

On the Artificial Culture of Marine Plankton Organisms.

By

E. J. Allen, D.Sc.,

Director of Laboratories and Secretary to the Council of the Marine Biological Association,
and

E. W. Nelson,

Assistant Naturalist.

	PAGE
Introduction	422
I. Culture of Plankton Diatoms	423
A. Practical Culture Methods	423
1. Miquel's Method	423
2. Houghton Gill's Method	426
3. (a) Modification of Miquel's Method, "Miquel Sea-water"	427
(b) English Channel Water	429
(c) Tank-Water	430
(d) Animal-Charcoal Water	431
(e) Peroxide of Hydrogen Water	434
(f) Cultures in these Media	435
B. Experiments with a view to Determining the Conditions which underlie the successful Culture of Diatoms	436
Methods	437
The Sea-water employed	437
The Constituents of Miquel's Solutions	438
Animal Charcoal and Peroxide of Hydrogen	443
Reviving Exhausted Cultures	443
Silica	444
Organic Infusions	445
Artificial Sea-water	446
Alkalinity	447
Salinity	453
Light	453
Temperature	454
General Conclusions	455
II. Mixed Cultures	457
III. Notes on particular species of Diatoms, on their methods of Reproduction, and on other Algae occurring in Cultures	460
IV. Rearing of Marine Larvæ	464
Methods	464
<i>Echinus acutus</i>	466
<i>E. esculentus</i>	466
<i>E. millaris</i>	467
<i>Cucumaria saxicola</i>	467
<i>Pomatoceros triqueter</i>	468
<i>Chaetopterus variopedatus</i>	468
<i>Sabellaria alveolata</i>	469
<i>Archidoris tuberculata</i>	469

422 ON THE ARTIFICIAL CULTURE OF MARINE PLANKTON ORGANISMS.

Colanus Annularis
Echinus hybrid
Saccellina carolin

Summary of Method for Rearing Larvae
Bibliography

Introduction. The observations to be recorded in this Paper were commenced in March, 1905. They originated in an attempt to find a general method for rearing marine larval forms. Several investigators had previously succeeded in rearing Echinoderms, Molluscs, and Polychaetes from artificially fertilized eggs under laboratory conditions, but the process was generally difficult and the results more or less uncertain. The most promising method seemed to be that adopted by Caswell Grave (26), who was able to rear his larvae by feeding them on diatoms. Grave obtained his diatoms by placing sand, collected from the sea bottom, in aquaria and using such diatoms as developed from this material. All the methods, however, suffered from the uncertainty of not knowing what organisms were introduced into the aquaria in which the larvae were to be reared, either in the original sea-water or along with the food-supply.

It appeared, therefore, at an early stage of the work, worth while to make an attempt to carry out rearing experiments on a more definite and precise plan, to endeavour, in fact, to introduce the larvae to be reared into sterile sea-water, and to feed them with pure cultures of a suitable food. This was the ideal to be aimed at. As a matter of fact, it has seldom, if ever, been attained in practice; nevertheless a considerable measure of success has been achieved by working upon these lines, and during the course of the work innumerable problems relating to the physical conditions under which plankton organisms can best flourish have presented themselves. Some account of the experiments made may be of interest to other workers, although many of the problems raised are not yet solved, notwithstanding the fact that some 1500 cultural experiments have been under observation. It is rather with a view of stimulating other work upon similar lines, than of bringing forward conclusive results, that this paper is being published.

In the summer of 1907, Mr. E. W. Nelson became associated with the investigation, and since that date the experimental work has been carried out by him. The discussions in this paper of a more chemical character, particularly the section on alkalinity, are almost entirely the work of Mr. Nelson, and we have both had throughout the advantage of the constant advice and help of Mr. D. J. Matthews on all such matters.

ON THE ARTIFICIAL CULTURE

I. CULTURE

A. PRAC

1. *Miquel's Method.* At Plankton diatoms; and t Miquel (11) for fresh-wat succeed with marine-botte

The essential features diatoms, are as follows:—

Two solutions are prep

SOLUTION A.

Magnesium sulphate
Sodium chloride
Sodium sulphate
Ammonium nitrate
Potassium nitrate
Sodium nitrate
Potassium bromide
Potassium iodide
Water

SOLUTION B.*

Sodium phosphate
Calcium chloride
Hydrochloric acid
Ferric chloride
Water

Forty drops of Sol added to each 1000 cc keeping it at 70° C. fo

According to Miqu material in the form Zostera. Maceration time before they are and sterilized. Orga

* "The preparation of as follows: To the Sodium of Hydrochloric acid, the Calcium chloride dissolve I call Phospho-ferro-calc throws down a slight broxide, which should be c⁺ "Acid chlorhydriq⁺ "Perchlorure de fer

I. CULTURE OF PLANKTON DIATOMS.

A. PRACTICAL CULTURE METHODS.

1. *Miquel's Method.* Attention was first directed to the culture of Plankton diatoms; and the methods, which had been elaborated by Miquel (11) for fresh-water diatoms and had been found by him to succeed with marine-bottom diatoms, were tried.

The essential features of Miquel's method, as applied to marine diatoms, are as follows:—

Two solutions are prepared:—

SOLUTION A.

Magnesium sulphate	10	gram.
Sodium chloride	10	"
Sodium sulphate	5	"
Ammonium nitrate	1	"
Potassium nitrate	2	"
Sodium nitrate	2	"
Potassium bromide	0.2	"
Potassium iodide	0.1	"
Water	100	"

SOLUTION B.*

Sodium phosphate	4	gram.
Calcium chloride (dry)	4	"
Hydrochloric acid	† 2	cc.
Ferric chloride	‡ 2	"
Water	80	"

Forty drops of Solution A and 10 to 20 drops of Solution B are added to each 1000 cc. of sea-water, and the sea-water is sterilized by keeping it at 70° C. for about 20 minutes.

According to Miquel it is also necessary to add "organic nutritive material in the form of bran, straw, or filaments of weed such as *Zostera*. Macerations of these should be made up separately, some time before they are required for use, and should be carefully filtered and sterilized. Organic matter must, however, be used very sparingly,

* "The preparation of Solution A presents no difficulty; Solution B should be made up as follows: To the Sodium phosphate dissolved in 40 cc. of water are added first the 2 cc. of Hydrochloric acid, then the 2 cc. of hydrous Ferric chloride and then the 4 gram. of Calcium chloride dissolved in 40 cc. of water, taking care to shake the mixture, which I call Phospho-ferro-calcic solution. The addition of this last solution to the maceration throws down a slight brownish flocculent precipitate, formed for the most part of Ferric oxide, which should be carefully separated from the liquid used for cultivations."

† "Acid chlorhydrique pur à 22°." Presumably meaning degrees Baumé = sp. gr. 1.169.

‡ "Perchlorure de fer liquide à 45°." As above = sp. gr. 1.421.

or else putrefaction will set in and the cultures will be irrevocably lost." As a matter of fact, we have found that such organic infusions are unnecessary, when dealing with plankton diatoms, and it has not been our practice to employ them (cf., however, p. 445).

Miquel obtained cultures of single species of diatoms either by picking out individual diatoms under the microscope and introducing them into the prepared water, or by adding a small quantity of water containing a mixture of diatoms and other organisms to some prepared water, and subdividing this into a number of tubes. If the subdivision has been carried out sufficiently some of the tubes may contain one kind of diatom only, from which fresh cultures can be made. In this way, by repeated subdivision, cultures can be obtained which, by inoculating fresh quantities of prepared water from time to time, may, with care, be maintained indefinitely. Such cultures, however, must practically always contain bacteria, and Miquel distinguishes them from bacteria-free cultures, which he terms "*Cultures des Diatomées à l'état de pureté absolue*." The latter he found very difficult to obtain, but, through repeated washing in sterile water, followed by fractional subdivision, he succeeded in getting some in which he could find no trace of bacteria by ordinary bacteriological methods (cf. Miquel 11, p. 155; cf. also Richter, 16-18).

We propose to call any diatom culture, which can be carried on practically indefinitely by inoculating fresh supplies of prepared water, a "*persistent*" culture, the term "*pure*" culture being reserved for cultures which can be proved to contain not more than one organism. We are not satisfied that we have yet succeeded in obtaining cultures of the latter kind. For the most part our persistent cultures contain one species of diatom only, and are free from all organisms larger than small flagellates.

In our earlier experiments with plankton diatoms, we obtained persistent cultures, containing a single species of diatom, by both of the methods recommended by Miquel. We, however, have rarely succeeded by picking out single diatoms or chains of diatoms, for although we have passed the selected diatom through several changes of sterilized sea-water, the resulting cultures, even when the diatoms have multiplied to some extent, have generally shown evidence of contamination by harmful organisms, and have soon died down. Only in one of the earliest experiments, and in one more recent, has complete success resulted. In the first case a small chain of six or eight frustules of *Skeletonema costatum*, picked out in April, 1905, gave rise to a culture which still persists (Nov., 1909). Subcultures can still be obtained even from the original flask inoculated in April, 1905. In the

second case a chain from a Petri dish.

The method of and persistent culture this way.

A more ready way to add one or two culture medium, dishes). The diatom from vibration, a *in situ*. The ten the dishes exposed being avoided.

different species of the Petri dishes and transferred colonies should a stage as possible a diatom or a whole of the water persistent culture above method is placed gelatinous successful persistence, which will from a very sterile sea-water objective content various species sea-water, were diluted plankton respectively persistent culture we may conclude short chains culture medium

We have succeeded in obtaining diatoms in p

second case a chain of 8 or 9 cells of *Chaetoceros densum*, picked out from a Petri dish culture, has given a particularly good growth.

The method of dilution and subdivision has been more successful and persistent cultures of a number of species have been obtained in this way.

A more ready method of obtaining the cultures is, we have found, to add one or two drops of plankton to, say, 250 cc. of a suitable sterile culture medium, and to pour this into shallow glass dishes (Petri dishes). The dishes should be placed in a position as free as possible from vibration, and where they can be easily examined with a lens *in situ*. The temperature should be kept as constant as possible and the dishes exposed to light of moderate intensity, direct sunlight being avoided. In the course of a few days, colonies of diatoms of different species will be seen at different spots on the bottom of the Petri dishes. These can be picked out with a fine pipette and transferred to flasks containing fresh culture medium. The colonies should be picked out from the Petri dishes at as early a stage as possible, because if left too long some one organism, a diatom or a flagellate, may have multiplied so rapidly that the whole of the water in the dish becomes infected with it. In this case persistent cultures of a single species would not be obtained. The above method is similar to one described by Miquel, excepting that he placed gelatinous silica at the bottom of the vessel. Some very successful persistent cultures were obtained from the following experiment, which will serve to illustrate the method:—A sample of plankton, from a very fine-mesh bolting-silk tow-net, was diluted down with sterile sea-water, until a single drop examined under a two-thirds-inch objective contained on an average ten organisms, chiefly diatoms of various species. Petri dishes (4 in.), containing 60 cc. each of Miquel sea-water, were then inoculated with various numbers of drops of the diluted plankton. The two dishes, to which two and three drops respectively were added, gave the best results; and from these persistent cultures of several species of diatoms were obtained. Hence we may conclude that the most advantageous number of single cells or short chains of cells to be added to a 4 in. Petri dish, containing 60 cc. culture medium, is about 20 to 30.

We have succeeded in obtaining the following species of Plankton diatoms in persistent cultures:—

Asterionella japonica, Cleve.

Biddulphia mobiliensis (Bail.), Grun.

Biddulphia regia (M. Schultze).*

* See pp. 461.

Chaetoceros denatum, Cleve.
Chaetoceros decipiens, Cleve.
Chaetoceros constrictum, Grun.
Cocconeis scutellum, Ehr. var. *minutissima*, Grun.
Coscinodiscus eccentricus, Ehr.*
Coscinodiscus Granii, Gough.
Ditylimum Brightwellii (West), Grun.
Lauderia borealis, Grun.
Nitzschia closterium, W. Sm.
Nitzschia closterium, W. Sm., *forma minutissima*.
Nitzschia seriata, Cleve.
Rhizosolenia stouterfothii, H. Perag.
Skeletonema costatum (Grev.)
Streptotheca thamensis, Shrubbs.
Thalassiosira decipiens, Grun.*

It is hardly necessary to add that in dealing with these cultures, similar precautions to those used in bacteriological work must be taken, all vessels and instruments being carefully sterilized before they are brought into contact with the prepared sea-water. The cultures are best made in small, wide-mouthed flasks, which may be plugged with cotton wool, or simply covered with watch-glasses. The flasks should be kept at as uniform a temperature as possible (from 12°-17° C.) and should be exposed to strong daylight, direct sunlight being avoided. A flask should not be more than half filled with culture fluid, so that the surface exposed to the air may be large in proportion to the volume of fluid.

Other Methods. The addition of the solutions devised by Miquel to sea-water has in all cases given us good cultures of diatoms, and the method is certain in its action. We have, however, made numerous experiments by treating sea-water in other ways, with a view to finding out what are the best conditions under which plankton diatoms will grow, and of arriving at some explanation of the action of the different salts contained in Miquel's solutions.

2. *Houghton Gill's Method.* H. Houghton Gill (5), a contemporary of Miquel, made use of a culture medium not essentially different from that employed by the latter. Unfortunately he died before publishing his work, but an account of his principal results is given by Van Heurek. In his final method Houghton Gill made use of four distinct solutions, as follows:—

* See p. 460.

SOLUTION 1.

Crystallized sodium phosphate	2 gm.
Calcium chloride	4 "
Syrup of iron chloride	0.5 "
Strong hydrochloric acid	1 "
Water	100 "

SOLUTION 2.

Crystallized magnesium sulphate	4 "
" sodium sulphate	4 "
" potassium nitrate	4 "
Common salt (sodium chloride)	8 "
Potassium bromide	0.2 "
Potassium iodide	0.2 "
Water	100 "

SOLUTION 3.

Crystallized sodium carbonate	4 "
Water	100 "

SOLUTION 4.

Well-washed, precipitated calcium silicate	25 "
Water	75 "

All the salts employed must be chemically pure. Three cc. of each of these liquids are added to 1000 cc. of fresh water or sea-water (according to circumstances), and the whole sterilized. In his earlier work Houghton Gill added a sterilized infusion of grass or of diatoms, but it is not clear from the accounts whether this was still employed with the above solutions. We have obtained very good cultures with the above solutions, to which we did not add any organic infusion.

3 (a). *Modification of Miquel's method. "Miquel Sea-water."*

Since several of the components in Miquel's formula for solution A (p. 423) are obviously unnecessary, when sea-water is being used as the basis of the culture-medium, we adopted for our own work the following modifications:—After some preliminary experiments it was found, as would be expected from the composition of sea-water, that the only salts of value to the medium are the three nitrates, KNO_3 , NaNO_3 , NH_4NO_3 , and possibly KBr and KI . The omission of the two latter was soon found to make no difference. Experiments also showed that the formula for solution A could, without any appreciable detriment to results, be further simplified to the one salt, KNO_3 , or NaNO_3 , but not NH_4NO_3 . At first the amount of KNO_3 , dissolved in 100 cc. distilled water, used to make the modified solution A, was the same as the sum of the weights of the nitrates in

Miquel's own formula, viz. 5 grm. But later experiments showed that a considerably greater concentration of KNO_3 than this gave more lasting cultures; the strength of solution, and amount to be added to a litre of sea-water, in order to obtain the best results, being 2 cc., 2M, KNO_3 .

In the case of solution B no modification has been adopted, but it has been found that small variations in the amounts of the ingredients used do not affect the results. A convenient method for measuring the right amount of FeCl_3 is to warm the salt until it just melts in its own water of crystallization, and to pipette out 2 cc. with a previously warmed pipette. No temperature corrections need be considered. Also 2 cc. of the ordinary pure concentrated hydrochloric acid at room temperature will suffice.

Our own formula for preparing Miquel sea-water is now:—

SOLUTION A.*

Potassium nitrate	20.2 grm.	} = 2M KNO_3
Distilled water	100 "	

SOLUTION B.†

Sodium phosphate ($\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$)	4 "
Calcium chloride ($\text{CaCl}_2 \cdot 6\text{H}_2\text{O}$)	4 "
Ferric chloride (melted)	2 cc.
Hydrochloric acid (pure, concentrated)	2 cc.
Distilled water	80 cc.

To each 1000 cc. of sea-water‡ add 2 cc. solution A and 1 cc. solution B, and sterilize by heating to 70°C . When cool, decant off the clear liquids from the precipitate, which will have formed when solution B is added to the sea-water.

As a rule our cultures were made in 60 cc. of this medium, contained in short-necked, wide-mouthed flasks of 125 cc. capacity, so that the proportion of air-surface to volume of liquid was large.

The medium was found to give constantly satisfactory results. On inoculation from a persistent culture of such diatoms as *Thalassiosira*, *Skeletonema*, *Chaetoceras*, etc., a growth visible to the eye is obtained in about ten days, and then multiplication takes place very rapidly. In from three weeks to a month's time a very considerable growth will be seen making a brown, flocculent mass at the bottom and back of the vessel containing the culture.

* This strength has only been used in the most recent experiments; and solution A in this paper, unless otherwise stated, means the five per cent solution of KNO_3 .

† For preparing this solution see p. 423.

‡ "Miquel water" seems to succeed equally well, whether it is made by adding Miquel's solutions to "outside water" or to "tank water" (cf. p. 437).

In from the exhaustion and of sterilized number certain often, after a and the culture food-stuffs co possibly by n interesting in in this medium. The oldest made at th Although t any diatom healthy gro

In old c much defo chromatopl from a sin within a c results see

At the exhibited, frustules

(b) *E* number Plymouth shore, ha an avera year is f

If a culture to fift algae, fl lost. these inocula obtain 70°C . gave centre

In from three to four months the culture begins to show signs of exhaustion and the frustules lose colour, but they do not, as in the case of sterilized outside and tank water, completely die off. A great number certainly do die, but some remain in a resting condition, and often, after a period of six months or so, these begin to multiply again and the culture regains its former vigour. This is probably due to the food-stuffs contained in the dead frustules going into solution again, possibly by means of bacterial action. This periodicity in cultures is interesting in that it resembles what takes place in the ocean. Cultures in this medium will persist indefinitely, so far as our experience goes. The oldest culture in our possession is one of *Skeletonema costatum*, made at the very commencement of this work, dated April, 1905. Although the frustules in this culture are quite unrecognizable as any diatom now, on making a subculture in fresh Miquel a normal and healthy growth can always be obtained.

In old cultures the diatoms are nearly always found to be very much deformed, and often appear to be only a mass of broken-down chromatophores. Whether regeneration can be successfully obtained from a single chromatophore, which must presumably be contained within a cell-wall of some kind, has not been definitely decided, but results seem to point in this direction.

At the start of a culture a tendency to teratological forms is often exhibited, but when the growth is well advanced, the shape of the frustules is usually quite normal.

(b) *English Channel Water* ("Outside Water").—In a large number of our experiments sea-water brought in from outside the Plymouth breakwater, and therefore taken at some distance from the shore, has been used. This is referred to as "outside water." It has an average salinity of about 35.0‰, and the temperature range for the year is from 8° C. to 16° C.

If a sample of "outside water" is inoculated from a persistent culture of a plankton diatom, a small growth is obtained in from five to fifteen days. But soon minute bottom forms of diatoms, other algæ, flagellates, infusoria, etc., appear, and the inoculated species is lost. The total growth of any form is never large. If the growth of these foreign forms is prevented by sterilizing the water before inoculation, a considerably better growth of the plankton form is obtained. The water was, as a rule, sterilized by simply heating to 70° C., which temperature was found to be quite adequate. Boiling gave equally good results, but the former was preferred, as less concentration due to evaporation took place. Even under these conditions

no permanent culture can be obtained, the diatoms soon beginning to lose colour and getting into an exhausted condition. Death takes place in from two to three months after the culture has been started, and in many cases considerably sooner. Long before inability to start new cultures, the test of death, has been established, the valves appear on examination quite colourless and practically empty.

Samples of outside water, taken at times when the quantity of plankton was widely different, gave no appreciable variation in the results obtained by culture methods. It is, however, doubtful whether differences in the amounts of growth in cultures, proportional to the seasonal variation in the quantity of phytoplankton, would be sufficiently marked to be appreciable.

The total growth under cultural conditions, although small for a culture, is very much greater than any natural plankton that has come within our experience.

(c) *Tank-water*.—"Tank-water" or water taken from the supply of sea-water circulating through the tanks of the Aquarium at Plymouth, shows some striking and interesting differences from "outside water." This water is pumped up from the sea, just below the Laboratory, into two large, covered-in, settling reservoirs, with a capacity of 50,000 gallons each. Pumping is only done at high water, spring tides, so as to get the least contaminated water, and no water is pumped that does not show a specific gravity, measured with a hydrometer, of $\rho^{17.5} = 26.00$ ($S = 34.00$) or over. The water is allowed to settle for about a fortnight before being used for the general circulation.

The tanks themselves are made of slate and glass, and the pipes which convey the sea-water to them are of vulcanite, so that the water does not come in contact with metal, excepting in the pumps, which are of cast-iron. The two settling reservoirs are used alternately, for about a week each. From time to time, tide and water allowing, waste is replenished, and about twice a year each reservoir is emptied, cleaned out and refilled. The aquarium takes about 20,000 gallons, and this is in circulation with one of the two 50,000-gallon reservoirs. An estimate of the amount of life in the tanks of the aquarium must be exceedingly rough, but the intensity of the larger forms of life is far greater than anything met with in natural waters. About 500 fish and 2000 invertebrates, including all forms as large as an *Actinia equina*, might be somewhere near the mark. So it will be seen that the accumulation of excretory products must be a by no means negligible factor. The flora of the tanks is very restricted, and is chiefly composed of minute forms of algæ. Minute naviculoid diatoms,

Ectocarpus, *Cladophora* are the commonest. *Laminaria*, do not, a great number unfilled, are not represented.

As in the case inoculated from a minute forms, etc. The ultimate growth often considerably have been obtained found to be greater.

In cultures of a very great interest noted. The culture persists for a remains normal.

(d) *Animal-culture*. of purifying the known and inland places. a tank, which some powdered a good growth experiments and these have culture of diatoms.

Animal culture sold in two gallons experiments subsequently appears to grow animal character is prepared.

1. A question

* Analysis

Ectocarpus, *Cladophora*, *Enteromorpha*, *Vaucheria*, and unicellular algae are the commonest forms. The large seaweeds, such as *Fucus* and *Laminaria*, do not live long if introduced. Plankton diatoms, although a great number must be pumped up when the reservoirs are being filled, are not represented.

As in the case of outside water, a sample of "tank-water," inoculated from a persistent culture, will only give a very small growth, minute forms, etc., soon multiplying and choking out the plankton form. The ultimate growth of minute unicellular algae other than diatoms is often considerable, and many quite unknown and unidentified forms have been obtained. The total growth of vegetable forms is always found to be greater than in the case of outside water.

In cultures of plankton diatoms made with sterilized tank water, a very great improvement on outside sterilized water was always noted. The culture of the diatom used to inoculate this medium persists for a considerable period, and the colour of the frustules remains normal for two to three months.

(d) *Animal-Charcoal Water*.—The use of animal charcoal, as a means of purifying the water in small aquaria, has for a long time been known and practised by those who have kept such aquaria in inland places. At an early stage in our experiments, water from a tank, which was not in a satisfactory condition, was treated with some powdered animal charcoal and filtered. It was noticed that a good growth of diatoms took place in this water. Systematic experiments with the use of animal charcoal were then commenced, and these have resulted in a method of great value, both for the culture of diatoms and for the rearing of pelagic larvæ.

Animal charcoal is made by the carbonization of bones,* and is sold in two grades known as "pure" and "commercial." Our earlier experiments were all made with "pure" animal charcoal, but subsequently the "commercial" animal charcoal was largely used and appears to give equally good, if not better results. In both cases the animal charcoal is used in the powdered form. Animal-charcoal water is prepared as follows:—

1. A quantity of sea-water is sterilized by heating it in a flask to

* Analysis of Animal Charcoal, from Thorpe's *Dictionary of Applied Chemistry*:—

Carbon	10.51
Ca., Mg. phosphates, Ca. fluoride, etc.	80.21
Calcium carbonate	8.30
Other mineral matter	0.98
	100.00

side tubulure. This tube is connected, by means of rubber pressure tubing, to another glass tube leading down from the bottom of a small inverted bell-jar, placed some height above (in our case 14 feet, which gives a pressure of ca. 6 lbs. to the square inch inside the jar). A screw pinch-cock on this connection serves as a tap. The carboy containing the treated water stands just above the bell-jar, and is fitted with a tightly fitting rubber bung, through which two tubes pass. One is an ordinary syphon, the other the only air inlet into the carboy. This latter automatically keeps the level of the water in the bell-jar




FIG. 1.—Diagram of apparatus for preparing sterile sea-water by filtration, without contact with metal.

constant, by closing the air-inlet as soon as the water covers the end of the tube. When filtering water, the *modus operandi* is as follows:—

The carboy is filled with tank-water, treated, and allowed to settle as before. The Berkefeld candle,* bung, delivery tube, and connections (see Fig. 1) are sterilized by boiling for half an hour and fitted into place from within. (The delivery tube is shaped so that any drops of water, accidentally running down outside it, do not enter the vessel receiving the filtrate; and the jar should be large enough to allow the hand to fit the filter into place without much trouble.) The pinch-cock is closed and the syphon from the carboy started, which will automatically stop if the bung has been properly fitted. This should be watched to avoid accidents. The pinch-cock is then opened until the water rises in the jar well above the top of the candle, but still leaving some air space. The lid can now be fitted into place and screwed down. The tightness of this joint can be tested by pouring a little water into

* No. 5. Porcelain mount; length 8 ins., diameter 2 ins.

the crack round the lid, and observing if any bubbles are formed when the pinch-cock is opened. If all is right, no bubbles will be seen, and a good stream of water will flow out from the delivery tube. Our apparatus will filter about twenty litres an hour, and the filtrate is exceptionally bright and clear. The candle should be sterilized every three or four days that the apparatus is in use, to avoid indirect contamination by growths of organisms through the substance of the filter.* The water while passing through this apparatus only comes into contact with glass, earthenware, and rubber, the use of metal having been purposely avoided.

(c) *Peroxide of Hydrogen Water*. As it seemed probable that the action of animal charcoal was due to contact oxidation with the oxygen occluded in the charcoal, experiments were made to determine whether a similar effect could be produced by the use of hydrogen peroxide (H_2O_2). This was used in two ways. In the first method a sufficient quantity of H_2O_2 was added to the sea-water to ensure complete sterilization (1 cc. of H_2O_2 of 20 vols. strength per 1000 cc. of tank-water was found to be satisfactory), and the excess of H_2O_2 was decomposed by adding manganese dioxide. The water was then filtered through filter cloth, and the filtrate appeared to remain quite sterile. Good cultures of *Chaetoceras constrictum*, *Biddulphia mobiliensis*, and *Skeletonema costatum* were made in this water, which seemed to be as good as water treated by the animal-charcoal method.

The second way of using the peroxide of hydrogen was to start with water sterilized by heating to $70^\circ C.$, and to add to this H_2O_2 in small quantities at a time, until its presence could just be detected on testing the sea-water with permanganate of potash. In these circumstances, the first amounts of H_2O_2 are decomposed in the oxidation of organic substances in the water, and a very slight excess of H_2O_2 persists. For tank-water 1 cc. of 1 vol. H_2O_2 per 1000 cc. was found to give the best general effect. Cultures grown in water prepared in this way developed satisfactorily, being practically equal to those made in animal-charcoal water, but they became exhausted rather quickly.

The treatment of aquarium water with ozone was also tried, as this seems to offer a possibility of treating large quantities of water,† such as the whole bulk of water in an aquarium circulation, without very considerable expense. Experiments on a small scale, which we were able to make, unfortunately only with imperfect apparatus, showed

* See Bulloch and Craw., *Jour. of Hygiene*, VI, No. 3 (1906) p. 409.

† The use of ozonized air for the purification of fresh water for town-water supplies has been adopted in some localities. See Bridge, J. H. Paper read before Franklin Institute, reprinted in *English Mechanic* (1907), pp. 369 and 392.

that water treated than untreated sterilized by the of water which and bright.

(f) *Cultures* results, which describe the p set up with th culture of a t or *Chaetoceras*

A. "Outside Small swamp be larg

B. "Outside Slig

C. "Tank-San will li

D. "Tank A not b ance

E. "Outside B will

F. "Outside F bec

G. "Tank lon

H. "Tank field t

K. "C di

L. "T a b

that water treated with ozonized oxygen gave distinctly better cultures than untreated water. Although the sea-water was not absolutely sterilized by the treatment to which we actually subjected it, a sample of water which was visibly clouded with bacteria became quite clear and bright.

(f) *Cultures in these Media.* In order to make clear the different results, which are obtained by using these different waters, we will describe the probable result which would be got from a series of flasks set up with the following media, and each inoculated with a persistent culture of a true plankton diatom, such as *Thalassiosira*, *Skeletonema*, or *Chaetoceras*.

- A. "Outside water" untreated.
Small growth in from five to fifteen days, almost immediately swamped by growths of foreign forms; the latter, however, will never be large.
- B. "Outside water" sterilized.
Slightly larger growth, very soon becoming exhausted.
- C. "Tank-water" untreated.
Same result as in A, but growths will be much larger, healthier, and will last longer.
- D. "Tank-water" sterilized.
A fair growth of the inoculated species, but the total growth will not be as great as in C; the diatoms will retain their normal appearance for some time.
- E. "Outside water" + Miquel's solutions A and B, sterilized.
Best culture in series, both in quantity and quality. The diatoms will remain normal and healthy for a very long period.
- F. "Outside water" sterilized and treated with Animal Charcoal.
Fair growth, especially at first; diatoms will soon grow pale and become exhausted; better than D.
- G. "Tank-water" sterilized and treated with Animal Charcoal.
As F, only growth will be slightly greater and will last considerably longer. Third best in series.
- H. "Tank-water" treated with Animal Charcoal and filtered through Berkefeld filter.
This will usually be the second-best culture in the series, but the difference between this and G will only be slight.
- K. "Outside water" treated with H_2O_2 .
This will most resemble F, but will not be quite so good.
- L. "Tank-water" treated with H_2O_2 .
A distinct improvement over K. This medium is rather variable, and in some cases the growth obtained has been quite equal to F, if not better.

B. EXPERIMENTS WITH A VIEW TO DETERMINING THE CONDITIONS WHICH UNDERLIE THE SUCCESSFUL CULTURE OF DIATOMS.

The attempt to make cultures of diatoms for use as food, when rearing pelagic larvæ, led naturally to an effort to determine the best culture medium and the most favourable conditions for the rapid and continuous growth of diatoms. Before success can be attained in this direction exact knowledge as to the nature of the essential food-stuffs—and in fact as to the general physiology of the Diatomaceæ—is necessary.* Numerous experiments, extending over the last three years, have been carried out, with a view to obtaining such knowledge, and the results, though still by no means complete or conclusive, are perhaps worth recording.

A great difficulty which has to be met in carrying out such investigations on marine diatoms is caused by the fact that, when sea-water is used as a basis for the culture media, we are dealing with a solution of a very complex and very variable character, the exact nature of which it is extremely difficult to determine. The most direct method of research, namely, chemical analysis, has not proved of much service, owing to the uncertainty and in many cases impossibility of accurate determinations, in sea-water, of such minute quantities of substances as those upon which the growth of plankton diatoms has been found to depend.

We have had, therefore, to rely, for the most part, on the lengthy and tedious process of analysis by "trial and error," the experiments being largely conducted on lines suggested by Liebig's well-known "law of minimums" (Pfeffer, vol. i., p. 413). The ideal at which we aim is to find a culture medium, with artificially prepared sea-water as its basis, such that the absence, or diminution in quantity, of any one of its constituents would have a profound effect upon the growth of diatoms in it. Whether the conditions regulating growth in such a medium would be at all comparable to the natural conditions of life in the sea is a question that would have to be decided by experiment, but in any case this could be made a starting point for much more definite research than has yet been attempted. Up to the present time we have not, unfortunately, succeeded in finding such a culture medium. Throughout the work we have had very great difficulty, in spite of much care and many precautions, in obtaining consistent results. It may even happen that, in two flasks containing the same culture medium, inoculated with the same culture of diatom and standing side

* For general references to literature see Bibliography, especially Miquel (12), Schier (18).

by side, under exactly identical conditions, quite different degrees of growth must therefore be frequently reported, and must not be felt in any conclusions which the

It must be remembered, also, that the diatoms that we have used, having been taken from the sea, this fact has probably had some influence upon our attempts to obtain consistent results with success.

Methods. In carrying out the experiments, the procedure has been as follows. The sea-water, after being filtered through a section the procedure has been as follows. The procedure has been prepared from sterile sea-water, and the media have always been used. The culture media were prepared in liquid, in short-necked, wide-mouthed bottles, and a number of cultures were tested in each series, standing in a row together in such a way that the results could be compared as similar as possible. Controls were included in each series, so that the results could be compared by reference to the controls. The results from a persistent culture of diatoms were compared with those from a great majority of cases was compared the different media, and the results were found to be identical, and although on a small scale, a litre was made up, so that the results might be avoided. The results of a comparative series of cultures were compared when the basis of any two cultures was the same, and the amount of growth was compared by eye alone. Any difference in appearance was described here as appreciable. A few drops from each culture were examined microscopically, and the results compared.

The sea-water employed. The sea-water employed in the culture media has been as follows. A general description of the accurate chemical analysis and the methods to make clear many different chemical methods of sea

by side, under exactly identical conditions, as far as can be recognized, quite different degrees of growth will be observed. All experiments must therefore be frequently repeated before entire confidence can be felt in any conclusions which they seem to indicate.

It must be remembered, also, that in all the persistent cultures of diatoms that we have used, bacteria have probably been present, and this fact has probably had some influence on the result. Unfortunately our attempts to obtain absolutely pure cultures have not met with success.

Methods. In carrying out the experiments to be described in this section the procedure has been as follows:—All media have been prepared from sterile sea-water, and sterile vessels and instruments have always been used. The cultures have usually been made in 60 cc. of liquid, in short-necked, wide-mouthed flasks of 125 cc. capacity. When a number of cultures were to be compared, the flasks were kept standing in a row together in such a way as to keep the physical conditions as similar as possible. Control cultures in standard media were included in each series, so that results from different series could be compared by reference to the controls. The various media were inoculated from a persistent culture of a species of plankton diatom, which in the great majority of cases was *Thalassiosira decipiens* (p. 460). When preparing the different media the methods used were, as far as possible, identical, and although only about 60 cc. was needed for a culture, a litre was made up, so that errors due to measuring very minute quantities might be avoided. The media were all freshly prepared for each comparative series of cultures, the same sample of sea-water being used, when the basis of any two or more was the same. Comparative estimates of the amount of growth in the different cultures were made by eye alone. Any difference between amounts of growth that has been described here as appreciable has always been accompanied by a marked difference in appearance to the eye on holding the cultures up to the light. A few drops from each culture were also, from time to time, examined microscopically as a test of the quality and purity of the growth.

The sea-water employed. The sea-water employed as a basis for the culture media has been either (1) "outside water" or (2) "tank-water." A general description of these will be found on pp. 429-431. An accurate chemical analysis of both types of water would probably make clear many difficult points, but, as already pointed out, no chemical methods of sufficient delicacy have yet been devised.

We have seen that if we compare "tank-water," i.e. water from the closed circulation of the Plymouth Aquarium, with off-shore sea-water *in situ*, a most obvious difference is the much increased density of the larger forms of animal life in the former, combined with the almost complete absence of plant life. Hence the concentration of excretory products in the tank-water must be very much higher than in outside water. Other factors, such as increased bacterial action, artificial aeration, etc., in tank-water, must also be taken into account (cf. Vernon, 58; Smith, 56). There seems to be direct evidence to show that the concentration of nitrates, possibly due to the action of nitrifying bacteria on the products of excretion, such as urea, ammonia, etc., is considerably higher in the tank-water, and the presence of soluble organic matter, in concentrations never met with in the sea, can almost certainly be assumed. It is probably due to the presence of these nitrates and soluble organic substances that sterilized tank-water is a much better medium in which to grow diatoms than sterilized outside water (see p. 435).

The constituents of Miquel's solutions. It has already been stated that no better medium for the culture of plankton diatoms has been found by us than the solutions recommended by Miquel, although these solutions may be modified and simplified in various ways with equally good results. The formulæ recommended by Houghton Gill give very similar cultures. The essential features of Miquel's and Houghton Gill's methods, when adapted to sea-water, are the same. Miquel's solution A, and Gill's solution 2, can both be replaced by a solution of potassium nitrate (p. 427). Again, Miquel's solution B and Gill's solution 1 only differ in the proportionate amounts in which the various constituents are prescribed. The formulæ are:—

	Miquel's sol. B.	H. Gill's sol. 1.
$\text{Na}_2\text{HPO}_4, 12\text{H}_2\text{O}$	4 gm.	2 gm.
Ca Cl_2	4 "	4 "
Fe Cl_3 (syrupus)	2 cc.	0.5 "
HCl (concentrated)	2 cc.	1 "
Water	80 cc.	100 "
	Use 1 cc. per 1000.	Use 3 cc. per 1000.

The proportionate amounts added to equal volumes of sea-water are:—

	Miquel's sol. B.	H. Gill's sol. 1.
Na_2HPO_4	10	12
Ca Cl_2	10	24
Fe Cl_3	5	3
HCl	5	6

Since culture using media prepared with Miquel's proportions of

We must note in Miquel's sea-water may be considered "protective,"* material must be used in the isolation of haptophytes and any more direct evidence not directly evidence

Our experiments with simple solutions and that the results are due to the amount

Inoculation

A. Normal

B. Diatoms

C. Diatoms

D. Diatoms

Consider the known action of that exhausts the nutrition of the assumption of the amount of a completely correct.

Consider the increased

* Loeb, † For this paper indicate Miquel's of each ad

NEW

Since cultures can be obtained with no appreciable difference by using media prepared by adding either of these solutions, together with Miquel's solution A, to sea-water, a considerable latitude in the proportions of the salts present is tolerated.

We must now consider what is the rôle of the various constituents in Miquel sea-water. The part played by any salt of a culture medium may be considered as being either, firstly, "nutritive," or secondly, "protective."* Under the first heading, any direct addition of food material must be included; under the second, any removal or neutralization of harmful substances, such as toxins and possibly bacteria, and any more remote effects, which, although influencing growth, do not directly enter into the metabolism of the plant.

Our experiments have proved that solution A can be reduced to a simple solution of potassium nitrate† without detriment (cf. p. 427), and that the amount of growth is, within limits, roughly proportional to the amount of KNO_3 added, as the following experiment shows:—

- Inoculated from persistent culture of *Thalassiostra decipiens*.
- A. Normal Miquel sea-water.
Growth as usual.
 - B. Ditto, but only one-half amount of sol. A.
Good growth at first, but exhausted sooner than A.
 - C. Ditto, but $2\frac{1}{2}$ times amount of sol. A.
Was slower than either A or B at start, but afterwards was better than A or B and lasted longer.
 - D. Ditto, but five times amount of sol. A.
As C, but in greater degree.

Considering the nature of the substance added, and its already well-known action in plant metabolism, these results, coupled with the fact that exhausted cultures can often be regenerated by the simple addition of nitrates (see below, p. 444), are quite consistent with the assumption that sol. A is simply nutritive in action. The concentration of nitrates in natural sea-water is so low (Brandt, 47) that the amount available in a culture of untreated water very soon becomes completely exhausted, and it is this deficiency that sol. A probably corrects.

Considering now the action of sol. B, it must first be observed that increased concentration of nitrates alone will not explain the whole

* :* Loeb, *The Dynamics of Living Matter* (New York, 1906), p. 77.

† For the sake of convenience, the expression sol. A will be used throughout the rest of this paper to indicate a simple solution of potassium nitrate (5 per cent) and sol. B to indicate Miquel's phospho-ferricalcic solution (p. 423). Unless otherwise stated, the amounts of each added to 1000 cc. sea-water will be normal, i.e. 2 cc. sol. A and 1 cc. sol. B.

An analysis of 1 cc. Miquel sol. B, the amount added to 1 litre Miquel sea-water, gave:—

P_2O_5		.00825	gm.
Fe_2O_3		.0105	"
CaO		.0145	"

Comparing these figures, it seems probable that, when added to sea-water, all the iron in sol. B is precipitated, and a certain amount also of the phosphate and calcium. The additive effect on the sea-water is therefore a slightly increased concentration of phosphate and calcium.

An analysis of a sample of tank-water for phosphorus, before and after treatment with sol. B (1 cc. per thousand), gave the following figures:—

Tank-water		.5	mgram. P per litre = .00163	gm. P_2O_5
Tank-water + sol. B				
(without precipitate)		1.5	" " " = .00488	" "

It will be noticed that the figures from the different analyses do not agree very well. This is probably due to the fact that different samples were used for analyses in each case, and also to the fact that the solutions were made up in the ordinary way, without any special precautions, volumes, for instance, being measured in cylindrical glasses, pipettes, etc.

Cultures were tried in sea-water containing the normal amount of sol. A, plus the normal constituents of sol. B, less all the iron and less the amount of phosphate that would combine with the iron to form basic ferric phosphate ($P_2O_5 \cdot 2Fe_2O_3 \cdot 12H_2O$). This solution should have very nearly the same chemical composition as normal Miquel sea-water from which the precipitate has been removed. Successful cultures could not, however, be obtained in it. Neither could cultures be grown in sea-water to which had been added the normal amount of sol. A and 1 mgram. P (as sodium phosphate) per litre.

To ascertain the effects of the different constituents of sol. B, experiments were carried out with separate solutions of these constituents, each of the same strength as in Miquel's formula. Different combinations of these solutions were added, together with sol. A, to sterilized sea-water, and the resulting media were inoculated in the usual way. It was found necessary to repeat these experiments a great number of times, as the results obtained were rather contradictory. To illustrate the methods used, a list of the different media, and notes on the cultures obtained in them, are given below. These media were inoculated from cultures of *Thalassiosira decipiens*, and the

cultures were kept under observation for at least four months. Series were made as uniformly as possible, and controls in standard media were included in each. The strength of the various solutions used in these experiments was the same as in Miquel's formula.

A. Outside water + sol. A + sol. B (normal Miquel sea-water).

First control.

B. Outside water + sol. A + Na_2HPO_4 sol. + FeCl_3 sol. + CaCl_2 sol.

Second control.

Good normal cultures were always obtained in these two controls.

C. Outside water + sol. A + Na_2HPO_4 sol.

A very uncertain medium. Sometimes no growth has been recorded and at other times a fair growth results, but these cultures are never equal to normal Miquel.

D. Outside water + sol. A + FeCl_3 sol.

Occasionally a very small growth has been obtained, but at the best it is very poor.

E. Outside water + sol. A + CaCl_2 sol.

About equal to D.

F. Outside water + sol. A + Na_2HPO_4 sol. + FeCl_3 sol.

Uncertain as C; no cultures have been obtained equal to the best in C.

G. Outside water + sol. A + Na_2HPO_4 sol. + CaCl_2 sol.

Some cultures very nearly equal to the controls have been obtained in this medium.

H. Outside water + sol. A + FeCl_3 sol. + CaCl_2 sol.

Poor, about equal to D.

Analysing the above results, we see that—

- (1) None of these modifications of sol. B give results equal to sol. B itself.
- (2) The best result is obtained from the combination of the phosphate and calcium chloride solutions.
- (3) Of the solutions used singly the phosphate is the best, the iron and calcium chloride being about equal.
- (4) The addition of FeCl_3 to Na_2HPO_4 , or the addition to CaCl_2 to FeCl_3 , does not improve the medium to any extent.

Experiments were also made to determine whether the precipitate thrown down in sea-water by Miquel's sol. B, itself had any influence on culture media. A quantity of this precipitate was prepared, filtered off, and then added to outside sea-water + sol. A (nitrates). A small growth was obtained, which was a distinct improvement on the control without the precipitate, but exhaustion soon set in.

Further discussion of the mode of action of sol. B, and as to whether that action is purely nutritive, or partly nutritive and partly

protective, is better post animal charcoal and oth

Animal Charcoal and culture medium for pl that prepared from a water gives at first al Miquel sea-water, bu appears much sooner, water," that is, tank-w powdered commercial filter. Tank-water as better than outside v concentration of nitrates

There is a very str charcoal and of Mi cultures, and the gre tank animal-charcoal sol. A is added to ani both in the colour a *Thalassiosira decipie* animal - charcoal w growth was obtained

It is possible the calcium, from the "nutritive" materi that its chief action gases, such gases l under normal cond

As was explained the action of ani parable to oxidati Fair growths of c the manner descr rapid exhaustion. water proved a m water.

Reviving Exhausted with water from

* Against this vie which has a still high coal, very poor cultur

protective, is better postponed until a later section, after the action of animal charcoal and other substances has been considered (see p. 455).

Animal Charcoal and Peroxide of Hydrogen. The most successful culture medium for plankton diatoms, next to Miquel sea-water, is that prepared from animal charcoal (cf. p. 435). Animal-charcoal water gives at first almost as good cultures of plankton diatoms as Miquel sea-water, but the tendency to paleness and exhaustion appears much sooner. The best cultures were obtained in "Berkefeld water," that is, tank-water from the Plymouth Aquarium treated with powdered commercial animal charcoal and filtered through a Berkefeld filter. Tank-water as a basis for animal-charcoal water is very much better than outside water, probably on account of the higher concentration of nitrates, etc.

There is a very striking resemblance between the effect of animal charcoal and of Miquel's sol. B upon sea-water used for diatom cultures, and the growths obtained by using tank-water + sol. B and tank animal-charcoal water are very similar in character. If Miquel's sol. A is added to animal-charcoal water, there is a great improvement, both in the colour and quantity of diatom growth, and in the case of *Thalassiosira decipiens* the chains are long and well formed. With animal-charcoal water + sol. B, on the other hand, practically no growth was obtained.

It is possible that a certain amount of phosphate, and perhaps of calcium, from the animal charcoal, goes into solution and serves as a "nutritive" material for the diatoms. But we are inclined to think that its chief action is "protective," and due to its power of occluding gases, such gases being in a state of higher chemical activity than under normal conditions.*

As was explained in a previous section (p. 434), the possibility that the action of animal charcoal might have some sort of effect comparable to oxidation, led us to experiment with hydrogen peroxide. Fair growths of diatom could be obtained in sea-water prepared in the manner described, but they always showed a tendency to rather rapid exhaustion. As in the case of animal-charcoal water, tank-water proved a much better basis for treatment with H_2O_2 than outside water.

Reviving Exhausted Cultures. Several experiments were carried out with water from old, exhausted cultures. The sediment was filtered

* Against this view would seem to be the fact, that when powdered cocoanut charcoal, which has a still higher power of occluding gases, was used in the place of animal charcoal, very poor cultures were obtained.

off, the filtrate was sterilized by heat, and then treated by various methods.

In one typical experiment the following was the result:—

Water from an exhausted culture of *Skeletonema costatum* in Miquel sea-water, reinoculated with the same diatom:—

- A. Filtered and sterilized.
No growth obtained.
- B. Ditto + sol. A (nitrates only).
Good culture, but did not last very long; further addition of nitrates made no improvement.
- C. Ditto + sol. B.
No growth.
- D. Ditto + sol. A + sol. B.
Very good growth, lasting considerably longer than B.
- E. Ditto + an. char.
No growth.

Exhausted cultures in animal-charcoal water gave the same general results on treatment and reinoculation. In an old culture of *Biddulphia mobiliensis* in outside water + sol. B only, which was in a very exhausted condition (nine months old), the addition of KNO_3 gave a very rapid regeneration, and the diatoms became of normal colour and form. This renewed growth, however, did not last very long, and a further addition of KNO_3 did not give any result. The addition of sodium phosphate also failed to stimulate growth. The same rapid regeneration, on the addition of potassium nitrate, has been obtained with almost every medium, but a second attempt has always failed.

Silica. A very noticeable character of the true plankton species of marine diatom is, that their skeletons are very markedly less silicious than the great majority of other forms. Their valves are only feebly marked, if at all, and they will not stand the vigorous treatment of cleaning with acids and heat that is commonly used in the case of fresh-water diatoms. In cultural forms, this absence of silica is still more obvious, and no marking can usually be seen on even those forms which, under natural conditions, are the most silicious, e.g. *Coscinodiscus eccentricus*. Deformed and distorted frustules are the rule in certain stages of growth in our cultures, and it is often very hard to make out more than the thinnest coating of silica. It is quite probable that this deformity can be accounted for simply by the absence of a strong silicious skeleton. As a rule, the more rapid the growth, the more teratological forms will be found. In untreated outside water little deformity will take place, but in normal Miquel, where very rapid growth takes place, the diatoms may assume almost

any conceivable shape to the normal age, the majority will be found that the of potassium silicate led to the collapse of the flasks in which possible that the demand, although cannot absorb silica is abundant. But or $\text{K}_2\text{Si}_2\text{O}_5$ for the We tried the acid in a culture of calcium silicate valves were much a trace of purple proved regular not seem to be exhausted culture

Organic Infusions. Infusions of culture in addition to if he ever did certainly state material is not of plants, the time, excellent fusions, both About a square an hour, could *Coscinodiscus* another, was

Infusions gave the same the final result out, these bacteria, in (7), in solution (Kutz), W. On placing

any conceivable shape. The form of the frustules tends to come back to the normal again, when the culture is well started, and in old stages the majority will be perfectly formed, although small and pale. It was found that the addition of silica (in early experiments as fragments of potassium silicate) was, as far as could be judged, immaterial, which fact led to the conclusion that a sufficiency dissolved out from the glass flasks in which the cultures were kept. During rapid growth, it is possible that the silica does not dissolve out fast enough to supply the demand, although it is also possible that diatoms, during rapid division, cannot absorb silica and form a perfect skeleton, even when the supply is abundant. Richter (18) has proved the necessity of either CaSi_2O_5 or $\text{K}_2\text{Si}_2\text{O}_5$ for the growth of *Nitzschia palea*, grown in pure cultures. We tried the addition of silica in various forms, and in one instance, in a culture of *Coscinodiscus eccentricus*, to which a little precipitated calcium silicate had been added, the uniformity and markings of the valves were much more regular than in the control. The presence of a trace of pure, dialysed silica, also, in one experiment, gave an improved regularity of form, but the quantity or rapidity of growth did not seem to be affected. No sign of regeneration could be obtained in exhausted cultures by the addition of silica.

Organic Infusions.—Miquel recommends the use in culture media of infusions of organic substances such as bran, straw, diatom broth, etc., in addition to the saline solution. He does not make it quite clear if he ever dispensed with them at all. In his general directions, he certainly states that the addition of both saline and organic nutrient material is necessary. As would be expected from the general metabolism of plants, the saline constituents are sufficient for growth. At the same time, excellent cultures have been obtained from dilute organic infusions, both with and without the addition of Miquel's sols. A and B. About a square inch of *Ulva* was boiled in 600 cc. sea-water for half an hour, cooled and filtered. In this medium an excellent growth of *Coscinodiscus eccentricus* in one case, and *Biddulphia mobiliensis* in another, was obtained, the growth lasting for some considerable time.

Infusions, made in the same way from a small piece of fresh fish, gave the same results, and although growth was rather slower at first, the final result was, if anything, slightly better. As Miquel points out, these infusions must be made very dilute, otherwise growths of bacteria, moulds, etc., will completely swamp the diatoms. Karsten (7), in some interesting experiments, showed that *Nitzschia palea* (Kutz), W.Sm., could be made to alter completely its mode of nutrition. On placing this diatom in organic nutrient solutions, it lost all chloro-

phyll and became colourless, but in saline media the chlorophyll would not regenerate, and the nutrition change back from heterotrophic to autotrophic.*

Of course, with our infusions, it cannot be said that the diatoms were necessarily feeding on dissolved organic material, as some necessary, saline, nutritive materials could have dissolved out from the weed or fish. If the former is the case, it might explain the superiority of tank-water over outside water, since the tank-water must contain a much higher percentage of organic substances in solution. If an alternative mode of nutrition autotrophic or mixotrophic could be proved, especially in the case of the "bottom" forms of diatoms, a great many phenomena could be explained, but the evidence is as yet far too slight to warrant any such assumption.

Artificial Sea-water.—As we have explained in a previous section, the ideal aimed at, in this part of our work, has been to obtain strong growths of diatomaceae in purely artificially prepared solutions of simple salts. If this end could be satisfactorily attained, the difficulties due to the unknown and variable composition of natural sea-water at once disappear. According to van 't Hoff (35) sea-water is a solution containing salts in the following molecular concentrations:—

NaCl 100.0, KCl 2.2, $MgCl_2$ 7.8, $MgSO_4$ 3.8, $CaCl_2$ 1.0 (varies).

Using these molecular concentrations, a sea-water of any desired salinity can be prepared. The chlorine content of average Atlantic water is about $Cl = 19.4$, and samples of artificial sea-water were prepared with the same chlorine value, thus:—

NaCl	26.75
KCl	.75
$MgCl_2$	3.42
$CaCl_2$	.51
$MgSO_4$	2.10
Double-distilled water	966.47
	1000.00

To make this solution comparable to natural sea-water, the "alkalinity" must be raised by the addition of an alkali such as Na_2CO_3 . After the importance of "alkalinity" as a factor had come before our notice, 2.4 cc. $M/2$ Na_2CO_3 was always added to the above solution in order to make the amount of base in equilibrium with CO_2 equivalent to the usual 40 mgrm. $OH^\circ/_{100}$ (p. 450).

* Cf. Zürnstein, *Zur Morphologie u. Physiologie d. Euglena gracilis*. Leipzig. 1899.

The only success we attained with artificial culture media was with four isolated experiments. Two of these were cultures in artificial sea-water + Miquel's solution were identical except that one was in flask and the other in a "resistance" between these two could be seen. The every way equal to normal Miquel sea over two years old. The other two of the same diatom in the same medium infusion, made by boiling up a small water. These gave just as good results factors from the weed detracts from of frequent attempts, over fifty in repeat this experiment, which may impurity in the salts or distilled media were prepared.

Alkalinity. Tornøe (43) and investigate the fact that sea-water excess of base over acid, which of sea-water." Dittmar defines measure of its potential carbonic his supposition that this excess dissolved CO_2 to form carbonate proportion, bicarbonates, is liable state of equilibrium actually (34) has shown, "sea-water actually just slightly more to the fact that sea-water of dissolved CO_2 .


If a salt solution with n OH° ions in concentration for pure water, be exposed to amount, depending on pressure into solution. This CO_2 weak acid H_2CO_3 , which free H° ions thus:—

The second stage of dissociation concentration of H° be

The only success we attained with artificial sea-water as a basis for culture media was with four isolated cultures in one of our earlier experiments. Two of these were cultures of *Coscinodiscus eccentricus* in artificial sea-water + Miquel's solutions A and B. The two cultures were identical except that one was in an ordinary Bohemian glass flask and the other in a "resistance glass" flask. No difference between these two could be seen. The growth obtained in both was in every way equal to normal Miquel sea-water, and is still fair, although over two years old. The other two successful cultures were growths of the same diatom in the same media, plus a small quantity of weed infusion, made by boiling up a small piece of *Ulva* in artificial sea-water. These gave just as good results, but the addition of unknown factors from the weed detracts from their general interest. In spite of frequent attempts, over fifty in number, we have not been able to repeat this experiment, which may possibly be due to some accidental impurity in the salts or distilled water from which the successful media were prepared.

Alkalinity. Tornøe (43) and Dittmar (33) were the first to investigate the fact that sea-water showed on analysis an apparent excess of base over acid, which excess they termed "the alkalinity of sea-water." Dittmar defines the alkalinity of sea-water as "a measure of its potential carbonate of lime," but this definition, and his supposition that this excess of base combines directly with dissolved CO_2 to form carbonates and, further, but only in very small proportion, bicarbonates, is liable to give a quite erroneous idea of the state of equilibrium actually occurring in the ocean. For, as Fox (34) has shown, "sea-water reacts *in situ* very nearly neutral, and actually just slightly more *acid* than distilled water." This is due to the fact that sea-water always contains a considerable quantity of dissolved CO_2 .

If a salt solution with *neutral* reaction, that is containing H^+ and OH' ions in concentrations equal to one another and the same as for pure water, be exposed to an atmosphere containing CO_2 , a definite amount, depending on pressure, temperature, and salinity, would go into solution. This CO_2 would combine with water and form the very weak acid H_2CO_3 , which would ionize with the formation of the free H^+ ions thus:—



The second stage of dissociation is so small as to be negligible. The concentration of H^+ being now increased, and OH' decreased, the

of cultures of plankton, the various sample culture media. T by Törnøe and D N/500, by intention bottles. Two ac were connected t by gravity. All with tubes of s known alkalinit mgrm. OH⁻/100), all operations b stored in stopp but those that l fresh standards with one anothe for diluting th still, redistilled sulphuric acid.

When carry sample and st flasks with a b was determin standard were back titrating indicator and lent was dete in the same n used. Let N to standard a time of pipet X the requir

Since all c no correction

In spite o by this me standard we and occasio (42) and its notice, and

solution would have an acid reaction. The actual amount of CO₂ thus dissolved would always be small; for instance, a salt solution of strength Cl=20.00 (average Atlantic water Cl=19.4) will at 10°C. dissolve about .3 cc. CO₂ per litre from an atmosphere containing 3% CO₂ (about normal). But the ocean is found to contain very much greater quantities than this, 60 cc., or two hundred times this amount, being a not unusual figure for the total CO₂. The difference between this amount and the .3 cc. or so dissolved by the neutral salt solution, as above, is kept in equilibrium with the 3% CO₂ of the atmosphere, by the amount of "excess" base equivalent to the amount of acid neutralized when an acid such as HCl is added to sea-water in excess. If a solution identical with sea-water but absolutely free from CO₂ (a practical chemical impossibility) could be obtained, then there would be present an excess of base over acid, and consequently an excess of OH⁻ ions over H⁺ ions, and an alkaline reaction. On exposing such a solution to the atmosphere, CO₂ would go into solution, ionize, and the H⁺ ions thus set free would react with the OH⁻ ions, due to the excess base, to form water. And this reaction would continue to take place, on more CO₂ dissolving, until all the excess OH⁻ ions were neutralized, at which point the solution would react neutral. Now, as before with the neutral salt solution, a further small amount of CO₂ would go into solution, bringing the solution into equilibrium with the atmosphere, and the excess H⁺ ions thus formed would give an acid reaction. The final result would be a solution exactly identical with natural sea-water. The total CO₂ found in sea-water can be considered as existing in two parts: the larger part in equilibrium with free base, its amount depending on temperature, pressure, and alkalinity; the smaller in equilibrium with the partial pressure of CO₂ in the atmosphere, its amount depending on temperature, pressure, and salinity. Although sea-water *in situ* has an acid reaction, it still maintains the property of being able to neutralize a certain amount of any acid stronger than H₂CO₃, that is any acid which, on dissociation, forms a higher concentration of H⁺ ions; for the stronger acid will turn out the H₂CO₃ in equilibrium with the "excess base" and CO₂ will be evolved.

In consideration of these points a less confusing definition of the "alkalinity of sea-water" would perhaps be a measure of its potential capability of neutralizing a strong acid* with the evolution of CO₂. This can be conveniently expressed, as is usual, in mgrm. OH⁻/100. Some of our earlier experiments seemed to show that "alkalinity" was a factor of considerable importance for the successful growth

* Such as HCl, with a high degree of ionization.

of cultures of plankton diatoms; so an attempt was made to analyse the various samples of water both before and after treatment as culture media. The method adopted was a modification of that used by Törnøe and Dittmar. Solutions of NaOH and H_2SO_4 of strength $N/_{100}$ by intention, were made up and stored in 5-litre "aspirator" bottles. Two accurately graduated burettes standing side by side were connected to these by tubes, so that they could be readily filled by gravity. All air inlets to burettes and stock bottles were fitted with tubes of soda lime. A standard solution of Na_2CO_3 of exactly known alkalinity, approximately that of average sea-water (40.00 mgrm. $OH\%$), was prepared by diluting down from a $N/_{10}$ solution, all operations being performed by weighing. These standards were stored in stoppered bottles of the fairly insoluble dark green glass, but those that had been kept for any length of time were not trusted, fresh standards being prepared. On analysis these standards agreed with one another to well within .1 mgrm. $OH\%$. The water used for diluting the standards was distilled water from the laboratory still, redistilled in all-glass apparatus with potassium bichromate and sulphuric acid.

When carrying out an analysis, equal volumes (about 100 cc.) of sample and standard were measured out into Jena glass Erlenmeyer flasks with a Knudsen automatic pipette. The specific gravity of each was determined by weighing in a 25 cc. pycnometer. Sample and standard were then titrated by running in acid from the burette and back titrating with alkali, using a 1% alcoholic solution of aurine as an indicator and keeping the liquid boiling. The acid to alkali equivalent was determined by titrating a pipetteful of double-distilled water in the same manner. The mean of at least four readings was always used. Let N and n be number of burette divisions of alkali equivalent to standard and sample respectively, and D and d their density at the time of pipetting out. Then if A is the alkalinity of the standard and X the required alkalinity of sample:—

$$X = A \frac{Dn}{Nd}$$


Since all operations were carried out at the same room temperature, no corrections for temperature are necessary.

In spite of the greatest care consistent results could not be obtained by this method of analysis. A sample analysed against the same standard would sometimes give results varying as much as 0.5 mgrm. and occasionally 1.0 mgrm. $OH\%$. The work on indicators by Salm (42) and its application to this question has only recently come to our notice, and it is our intention to experiment on this in future research.

The figures quoted below as the results of analyses have been rounded off as whole numbers, since their interest lies in their comparative rather than their absolute value, for convenience they are quoted as "alkalinity," although we are fully conscious that the methods used do not warrant this assumption, and that their actual chemical significance is still obscure.

The mean value for "outside water" was found to be fairly constant at 40.0 mgrm. $\text{OH}^{\circ}/_{\text{cc}}$, which figure agrees with results obtained by others for average ocean water. Samples from the aquarium tanks never gave as high figures as this, the average being approximately 37.5 mgrm. $\text{OH}^{\circ}/_{\text{cc}}$. From this it seems that the amount of base in equilibrium with CO_2 in tank-water is appreciably less than in outside water. A series of thirteen samples taken from seven miles beyond the Eddystone to well inside the Cattewater (an inner tidal harbour near Plymouth) showed a gradual lowering of the alkalinity from the normal 40, to 38 mgrm. $\text{OH}^{\circ}/_{\text{cc}}$ as the water became more estuarine and polluted.

The addition of Miquel's sol. B to sea-water was found, on analysis, to reduce the "alkalinity" by an amount equivalent to 10 mgrm. $\text{OH}^{\circ}/_{\text{cc}}$ or more. The 1 cc. sol. B added to a litre of sea-water, in itself contains a certain amount of free acid, equivalent to less than 4 mgrm. $\text{OH}^{\circ}/_{\text{cc}}$. But this reduction of alkalinity cannot be accounted for by the addition of free acid alone, because if only a quarter of the amount of sol. B is added, the alkalinity of the sample will be found to be, if anything, only very slightly higher. Also, if the various constituents of sol. B are added as separate solutions, thus obviating any addition of free acid, a reduction equivalent to about 6 mgrm. $\text{OH}^{\circ}/_{\text{cc}}$ is still obtained. The presence of ferric chloride in sol. B gives a possible explanation of this phenomenon. If a solution of ferric chloride is added to a solution of a soluble carbonate, a reaction, which can be expressed by the following equation, takes place:—



When the ferric chloride is added to sea-water, the final result will be that a certain amount of the "excess base" which was in equilibrium with CO_2 , will then be in equilibrium with the chlorine, available on the precipitation of hydrated ferric oxide, with a consequent liberation of CO_2 , and a reduction in "alkalinity" will, therefore, take place.

An analogy between the actions of Miquel's sol. B and animal charcoal can be seen in the fact that water treated with animal charcoal also shows a reduced "alkalinity," the amount being very variable in different samples.

Sea-water treated with animal charcoal, but in a much less degree, were used.

Control experiments were made with sea-water treated with these substances. It was found in obtaining an equilibrium (which may be judged, distilled water) showed a negative alkalinity of $\text{OH}^{\circ}/_{\text{cc}}$, and in the case of sea-water equivalent to 6 mgrm. $\text{OH}^{\circ}/_{\text{cc}}$. These results are only approximate, as the above results are due to the fact that the cloaks the true alkalinity.

Before any attempt was made to obtain a considerable difference in the media had presented themselves, a result was found. In the case of NaHCO_3 , this result was found to be outside water + sol. B. In the case of NaHCO_3 , this result was obtained in either 1 cc. HCl (p. 10) added 1 cc. HCl (p. 10) alkalinity of the 1 cc. normal growths restate that a weak diatoms.

In our most recent experiments for alkalinity, and for determining the effect of determining the media made in the following table:

- | | |
|---------------|-----------|
| A. Tank-water | Poor |
| B. Tank-water | Very poor |
| C. Tank-water | Very poor |
| D. Tank-water | Fair |

The sample of sea-water gave very poor results, that any improvement

Sea-water treated with H_2O_2 also showed a lowering of the alkalinity, but in a much less degree when, as usual, minimal quantities were used.

Control experiments on double-distilled water, which had been treated with these substances, were tried, but great difficulty was found in obtaining an end point with the indicator. As far as could be judged, distilled water treated with sol. B (quantities as with sea-water) showed a negative "alkalinity," equivalent to about 8 mgrm. OH^{100} , and in the case of animal charcoal a positive alkalinity equivalent to 6 mgrm. OH^{100} , but the colour change was so slow that these results are only the roughest estimates. The possibility that the above results are due to some effect on the indicator, which entirely cloaks the true alkalinity, must always be taken into consideration.

Before any attempts at analysis had been made, the probability that considerable differences might be found in the alkalinity of the various media had presented itself. Improvement in the growth of diatom cultures was found to result from the purely empirical addition of $NaHCO_3$, this result being most marked in normal Miquel sea-water, outside water + sol. B only, and Berkefeld water. No growth could be obtained in either tank-water or Miquel sea-water to which had been added 1 cc. HCl (pure, concentrated) per litre, but on again raising the alkalinity of the latter by the addition of $NaHCO_3$ or KOH good normal growths resulted. Richter (18) and H. Gill (5), also, both state that a weak alkaline reaction is necessary for the growth of diatoms.

In our most recent experiments, all the media have been analysed for alkalinity, and those given in detail below illustrate the importance of determining this factor. Cultures of *Thalassiosira decipiens* were made in the following media:—

A. Tank-water. Control.

Poor growth, hardly normal. Later, good growth of minute forms, etc.

B. Tank-water treated with cold commercial animal charcoal and filtered.

Very good growth indeed.

C. Tank-water treated with cold pure an. char. and filtered.

Very poor growth, comparable to A without minute forms.

D. Tank-water treated with pure an. char. as in C, but the an. char. was added red hot.

Fair growth, much superior to C, but not up to B.

The sample of pure an. char. used here had been previously found to give very poor results, and it was also quite contrary to our experience that any improvement in growth should be obtained by adding it hot.

But if we examine the results of analyses of these media for alkalinity a probable explanation presents itself. The following figures are only comparative:—

A—38 mgrm.	OH [°] /...	(used as standard).
B—37 "	" "	(higher than usual).
C—16 "	" "	(very low indeed).
D—34 "	" "	

It will be seen that the amount of growth in each treated sample follows the alkalinity very closely.

Solutions of Na₂CO₃, NaHCO₃ and HCl were made up, so that 4 cc. of any one contained an amount of acid or alkali equivalent to 10 mgrm. OH. From these a series of normal Miquel sea-waters of different alkalinities were prepared. Cultures of *Thalassiosira decipiens* were grown in these media.

- A. Normal Miquel sea-water. Control. A = 32.7 mgrm. OH[°]/...
Perfectly normal growth.
- B. Ditto + 4 cc. Na₂CO₃ per litre. A = 41.7 mgrm. OH[°]/... (= +9.0).
No difference between this culture and A.
- C. Ditto + 8 cc. Na₂CO₃ per litre. A = 50.2 mgrm. OH[°]/... (= +17.5).
Best culture in series in quality and quantity.
- D. Ditto + 4 cc. NaHCO₃ per litre. A = 42.4 mgrm. OH[°]/... (= +9.7).
Slightly better than control.
- E. Ditto + 8 cc. NaHCO₃ per litre. A = 51.5 mgrm. OH[°]/... (= +18.8).
As D.
- F. Ditto + 4 cc. HCl per litre. A = 22.2 mgrm. OH[°]/... (= -10.5).
Fair growth but never up to control, exhausted much sooner.
- G. Ditto + 8 cc. HCl per litre. A = 11.1 mgrm. OH[°]/... (= -21.6).
Poorest in series.

Except in the cases where the alkalinity was lowered by the addition of HCl, the results obtained from this series were not up to expectation. Nevertheless the majority showed a distinct improvement from increased "alkalinity" and in C, where the alkalinity had been raised 17.5 mgrm. OH[°]/..., this improvement was very marked.

Another point illustrated by cultural experiment is that in two samples of an. char. water, one with "outside" and the other with "tank-water" as a basis, the amount of growth in the latter considerably exceeded that in the former, and at the same time it was found that, with the tank-water, the alkalinity had not been reduced to the same extent as in the case of the outside water.

How far apparently anomalous results, which have so frequently

* Figures in parentheses are difference in alkalinity from control, in mgrm. OH[°]/...

occurred in our experiments changes in "alkalinity," ca

Salinity.—The salinity of the outside sea-water) of the outside between small limits, S = is also fairly constant, though only pumped up into the unless the salinity on taken. Experiments taken on the growth of diatoms various salinities were with double-distilled water slow evaporation. The evaporated down to the tion. Miquel solution solution was divided distilled water was added to the last, so that a from normal to near amount of Miquel's mixed culture of *S. Coscinodiscus eccentricus* two with highest practically sterile a limit of concentration. In the same way cultures of the same 100% did not seem quality of growth. the cultures of low The range of salinity S = 60‰ and was observed, except in noted.

An attempt to solutions was tried took place, although ocultating a culture weeks gave no growth

Light. Of all

* i.e. from

occurred in our experimental work, could be explained by unforeseen changes in "alkalinity," can only be answered by future research.

Salinity.—The salinity (or amount of salts dissolved in a litre of sea-water) of the outside water used in these experiments only varied between small limits, $S = 34.5 - 35.5‰$. The salinity of "tank-water" is also fairly constant, the average being about $S = 34.9‰$; water is only pumped up into the reservoirs at high water, spring tides, and unless the salinity on analysis is well above $S = 34.5‰$, no water is taken. Experiments to show what effect salinity pure and simple had on the growth of diatoms were undertaken. Samples of sea-water of various salinities were prepared by diluting down "outside water" with double-distilled water, and by concentrating "outside water" by slow evaporation. Two litres of "outside water" $S = 34.9$ were evaporated down to the bulk of one litre, giving a 50%* concentration. Miquel solutions 4 cc. A, 2 cc. B, were now added, and the solution was divided into ten culture vessels, 20 cc. in each. Double-distilled water was added, 2 cc. to the first, 4 cc. to the second, 20 cc. to the last, so that a series of media were obtained, varying in salinity from normal to nearly 50% concentration, each containing the same amount of Miquel's nutrient solutions. These were inoculated from a mixed culture of *Skeletonema costatum*, *Biddulphia mobiliensis*, and *Coscinodiscus eccentricus*. A good growth took place in all except the two with highest concentration. Of these two, the last remained practically sterile and the growth in the other was very poor. The limit of concentration, therefore, seems to lie between 35 and 40%. In the same way series of lowered salinities were prepared, and cultures of the same diatoms were grown in these. Dilution up to 100% did not seem to make any difference at all in the quantity or quality of growth. In a series extending the dilution to 200% even in the cultures of lowest salinity, a fair quantity of growth took place. The range of salinities covered by the various series was $S = 12‰$ to $S = 60‰$, and within these limits no effect on growth could be observed, except in the very highest, where a distinct deterioration was noted.

An attempt to grow *Coscinodiscus eccentricus* in tap water + Miquel's solutions was tried, and it was thought that some slight multiplication took place, although it was certainly not at all considerable. Inoculating a culture of normal Miquel sea-water from this after six weeks gave no growth.

Light. Of all the factors controlling the rate of growth of a culture,

* i.e. from every 100 cc. sea-water 50 cc. H_2O had been subtracted.

light seems to be by far the most important. Without light a culture soon dies off completely, showing marked signs of malnutrition very soon after having been placed in the dark, the brown pigment being the first to go, and later the chlorophyll. A culture (*Thalassiosira*) placed in the dark for five months was found to be completely killed, the diatoms being quite colourless. In cultures kept in bulbous flasks or any spherical vessel, the strongest and earliest growth always takes place at the side of the vessel away from the source of light, where the light will be found to be concentrated owing to the lens effect of a sphere of water. By painting a flask black on the outside up to the water-line of the medium, a very marked diminution in the rate of growth was obtained. The total growth was not affected, but depends on the available quantity of food-stuffs present.

Experiments on the reaction of cultures to different rays of the spectrum, obtained by coloured glass, were tried, but no results obtained. Miquel obtained marked results with yellow light, but in our experience, with plankton diatoms, satisfactory cultures could not be obtained under these conditions.

Temperature. The highest temperature which diatoms and allied forms can stand was about uniform for all the species tested, and lay between 35°–40° C. Cultures of the following species, viz. *Asterionella japonica*, *Nitzschia closterium*, minute naviculoid diatom, *Pleurococcus mucosus*, *Chilomonas* sp., were slowly heated in a water bath, and at every rise of 5° C. from 15° C. to 45° C., a few drops of the culture were pipetted out and a fresh flask inoculated. In all the flasks cultures were obtained where the heating process had not been carried above 35° C., but none in those where the temperature had exceeded this.

In the earlier stages of experimentation the cultures of diatoms were kept in various places about the Laboratory, and so were under quite different temperature conditions. Those placed in the warmer situations, i.e. near hot-water pipes, as a rule gave the most satisfactory results. In all the later work the cultures have been kept in one room, and an attempt has been made to keep the temperature of this room as nearly as possible constant at 15° C. A continuous record of its temperature has been kept by means of a recording thermograph, and no very great change of temperature has been noted. In a few isolated cases the temperature has dropped as low as 9° C., and in hot weather has risen just above 20° C., but these have only been for very short periods, the average temperature having kept remarkably constant. An apparatus in which flasks could be kept at

different uniform temperature, was used, but no results. About 17° C. seemed to be below this temperature was

General Conclusions. The experiments described in determining the conditions for the growth of diatoms, may now be divided into two parts. The first involved the making of a medium, and the second many of the questions raised will be necessary before

If we wish to obtain a pure plankton diatom, the conditions must be different from all other organisms. A "persistent" culture, under heat or filtration, and under sterile conditions. In the culture medium, it seems that the addition of nitrates, and perhaps this simple addition. Some other action, such as charcoal, or by percolation, is necessary in nearly every case. It is not possible to conclude conclusively that the cultures were purely nutritive, but the amount of the ingredients in the amount of the growth, the quantity of growth, the modification introduced, and the sol. B can be modified to have an appreciable effect, but are exceeded, then by supplying a necessary amount of animal charcoal, but, since the addition is comparable with the phosphates, there is a change in composition, we are not sure of the effect, they must be the results obtained, "nutritive," although

different uniform temperatures from 10° to 25° C., by means of hot air, was used, but no really satisfactory result could be obtained. About 17° C. seemed to give the maximum growth, and the cultures below this temperature were usually superior to those above.

General Conclusions. The general conclusions to be drawn from the experiments described in this section, which were made with a view to determining the conditions that underlie the successful culture of diatoms, may now be discussed. Although the experiments have involved the making of some 750 different cultures, our conclusions on many of the questions raised are still indefinite, and much further work will be necessary before a satisfactory answer can be given to them.

If we wish to obtain the maximum quantity of healthy growth of a plankton diatom, the diatom must first be obtained as free as possible from all other organisms, if not in a "pure" culture, at least in a "persistent" culture. All culture media should be sterilized either by heat or filtration, and the experiments should be conducted under sterile conditions. Starting with normal sea-water as the basis for the culture medium, it seems to be first necessary to raise the concentration of the nitrates, and possibly also of the phosphates, in solution. But this simple addition of nutrient materials will not in itself suffice. Some other action, such as that exerted by Miquel's sol. B, by animal charcoal, or by peroxide of hydrogen, seems to be imperative in nearly every case. The exact nature of this action we have not been able conclusively to determine. If the substance contained in sol. B were purely nutritive in character, we should expect that, when alterations in the amounts of the different ingredients were made, or when any one of the ingredients was omitted altogether, the differences in the quantity of growth would show a direct relation to the kind of modification introduced. But our usual experience has been that sol. B can be modified within certain limits, without producing any appreciable effect upon the resulting cultures, whilst if these limits are exceeded, there is an almost complete inhibition of growth. In supplying a necessary increase of phosphates, both Miquel's sol. B and animal charcoal may and probably do act as "nutritive" substances, but, since the addition of phosphates alone does not yield cultures comparable with those produced by either of these, and since, excepting phosphates, there is no possible common nutritive substance in their composition, we are led to conclude that, in addition to any nutritive effect, they must exert some other action. This view is supported by the results obtained by using H_2O_2 . This substance cannot be directly "nutritive," although it may be so indirectly, by oxidizing into useful

food-material substances which the diatoms are incapable of using in their metabolism, e.g. nitrites into nitrates. The absence of any increase in phosphates, when using H_2O_2 , may possibly be the reason why better results were not obtained with this medium. The action which, in addition to any nutritive value, we must assume that sol. B, animal charcoal, and H_2O_2 can all effect, would appear to fall into the class of "protective" actions (p. 439). It is quite conceivable that, with different samples of sea-water, this "protective" action is not necessary in every case, and this would account for the anomalous results met with when using sea-water + nitrates + phosphates only, in which medium sometimes good cultures, but more often the reverse, are obtained. The effect of Miquel's sol. B, animal charcoal, and H_2O_2 on the "alkalinity" of the sea-water, also points to some chemical change, which does not directly enter into the metabolism of the plants.

It may be pointed out that the action of such substances as finely powdered carbon, and ferric oxide precipitates, has been shown to produce a favourable effect on nutrient solutions used for the culture of certain higher plants, and it has been suggested that the beneficial action of these substances is the removal of toxic elements from the media (Breazeale, 3). Such removal of toxins would fall under our definition of "protective" action.

Of nutritive substances, other than those already mentioned, we have still to consider (1) silica and (2) dissolved oxygen and carbonic acid. Having regard to the conditions under which our cultures have been grown, i.e. in glass flasks, the question of silica does not seem to enter into the problems which we have discussed. A few words must, however, be said as to the dissolved gases. Whipple (62) and Baldwin (44) have drawn attention to the observed relations, which are found in natural waters, between algal growths and the amounts of dissolved oxygen and carbonic acid. That these factors are of great importance cannot be doubted, but in our cultures it seems reasonable to suppose that the conditions of saturation of these gases are the same in all, since series of cultures in standard media, such as Miquel sea-water or Berkefeld water, can be set up with the certainty that, if not every one, at least a very high percentage, will give normal results.

Of the purely physical factors, light is by far the most important. Within limits, the rate of growth in a suitable medium seems to depend directly on the intensity of the light (Whipple, 60). Absence of light, as would be expected, soon completely kills the diatoms.

Temperature also seems to affect the rate of growth to a certain extent, but for those temperatures at which we have experimented it does not appear to alter the quantity of growth.

Salinity, apart from the fact that it can be varied, does not appear to affect the growth of diatoms.

In what has been said, the persistent culture of diatoms is compared, comparatively, with the study of other organisms. The study of diatoms is not without interest.

A number of diatoms, about 10,000, were placed in a tall jar in constant motion. The water was changed, and a cover net (150 meshes) was used. A rich sample of diatoms was spring and summer the same, with the same nature of the water.

During the summer the rapid multiplication of diatoms was not always the case. They died off, though plankton diatoms during the early part of the summer in the body of the water. At the end of a summer, seen in every jar with the *Skeletonema*, *Lauderia borealis*, *Ditylisma*, *decipiens*. They only, when they died down to the bottom of the jar, flagellates of plankton diatoms experiments.

Salinity, apart from the quantities of available nutrient materials, can be varied within large limits without appreciable effect on the diatoms.

II. MIXED CULTURES.

In what has been said up to the present, we have been dealing with persistent cultures containing a single species of diatom, which are comparatively, if not entirely, free from admixture of other organisms. The study of cultures which contain a considerable mixture of organisms is not without interest.

A number of experiments have been made on the following lines. About 10,000 cc. of water, taken at some distance from shore, was placed in a tall bell-jar fitted with a "plunger," which keeps the water in constant movement. (*Journ. Mar. Biol. Assoc.*, Vol. 5, p. 176). The water was treated with Miquel's solutions in normal proportions, and a considerable quantity of plankton taken with a fine-meshed net (150 meshes to the inch) was added, say, 10 or 20 cc. of a moderately rich sample of tow-netting. The experiments were made during the spring and summer months, and the general course of events has been the same, with a certain amount of difference in detail according to the nature of the plankton present at the time.

During the first two days the water often became cloudy, owing to the rapid multiplication of small flagellate infusoria, though this was not always the case. Plankton Copepods and other animals gradually died off, though some survived for as long as a week or ten days. The plankton diatoms, on the other hand, generally multiplied rapidly during the early days of the experiments, the first to become abundant in the body of the water being usually *Skeletonema costatum*, which at the end of a week might be so thick, that a number of chains could be seen in every drop of water examined with the microscope. Along with the *Skeletonema* were found other plankton diatoms, such as *Lauderia borealis*, *Chaetoceras* (two or three species), *Biddulphia mobiliensis*, *Ditylium Brightwellii*, and in nearly every case *Thalassiosira decipiens*. These latter diatoms were present in moderate numbers only, when the *Skeletonema* was at its height; but as the *Skeletonema* died down they increased in quantity. At the same time *Nitzschia closterium* commenced to appear, both amongst the precipitate on the bottom of the jar, and in the general body of the water. Small green flagellates often began to get numerous also at this stage. The true plankton diatoms were at their height about a fortnight after the experiments were started. At this time a great many diatoms of all

kinds were to be found amongst the precipitate at the bottom of the jar, *Asterionella japonica* and *Coscinodiscus eccentricus* being often numerous here. During the course of the next week, however, *Nitzschia closterium* rapidly increased in quantity until, not only the sides of the jar were coated with it, but the whole mass of the water became thick and opaque. By this time the plankton diatoms had all disappeared, with the exception of those which may survive for a considerable period amongst the precipitate at the bottom of the jar. Bottom diatoms (*Navicula*, etc.) had begun to grow on the sides of the jar, and small green and brown algae (*Pleurococcus mucosus*, *Ectocarpus*, etc.) also appeared. Infusoria (*Euplotes* and other smaller forms) then became numerous, and as the *Nitzschia* and bottom diatoms increased on the glass, large numbers of *Amoeba* made their appearance among them. The jars continued in this condition for many months, the algae becoming more and more predominant.

From these experiments, as well as from instances of mixed cultures obtained in the course of our attempts to secure persistent cultures of single species of diatoms, it seems usual that, in a culture obtained by inoculating Miquel sea-water with plankton taken freshly from the sea, the true plankton diatoms are the first to develop in considerable numbers. Subsequently bottom diatoms and algae of various kinds become abundant, and the true plankton forms die out.

A complete explanation of this sequence of events would probably be of a very complicated character, and we have practically no evidence from our experiments which bears very directly on the question. It would seem, however, that the early predominance of the plankton forms in the cultures would naturally follow from the fact that, in the plankton material used for inoculation, these plankton forms are numerous, whilst bottom diatoms and spores of algae are rare. The subsequent very great predominance of such a species as *Nitzschia closterium* may be due simply to a very much more rapid growth rate, though it is difficult to avoid the impression that the organisms which finally take possession of the cultures, are in some way directly inimical to those which they supersede, not merely by robbing them of their food-supply, but perhaps, also, by the production of toxic substances. This suggestion does not, however, give an adequate explanation of the essential facts concerning these organisms. We have to consider two sets of species: (1) the true Plankton forms, which flourish in the open sea and can be grown quite easily in the laboratory, provided the cultures remain pure; and (2) what we may call "aquarium" or "bottom forms," which under experimental conditions invariably take possession, when present in mixed cultures, whilst the plankton forms

are killed off. Why is it always present in small numbers? Are there altogether other conditions such as succeed in gaining the difference between the sea and in the culture? A fruitful field for further research is so slight that further service.

The details of two subject of mixed cult

A flask, containing solutions, was inoculated with persistent cultures of following diatoms were *Biddulphia mobilis*, *Streptotheca thamensis*, introduced, since it was a new experiment was started on *Biddulphia*, *Coscinodiscus* were very healthy. was found.

On October 2nd *Coscinodiscus* was 1 and *Chaetoceros* were very numerous.

On October 3rd condition, *Coscinodiscus* flagellates (*Chlorococcum*) red colour.

Subsequently having probably increased very inconspicuous.

On July 28th were still in a h the flask. A g was also present subculture was subculture gave however, of a

are killed off. Why is it that, although species of the second class are always present in small numbers in plankton taken from the sea, they are there altogether outnumbered by the true Plankton forms, whereas under conditions such as those of our experiments they invariably succeed in gaining the upper hand? What are the factors which determine the difference in behaviour of these two sets of organisms in the sea and in the culture vessels? The whole question offers a very fruitful field for further experiment. The evidence at present available is so slight that further discussion of it here is not likely to be of much service.

The details of two experiments which we have made, bearing on the subject of mixed cultures, may, however, be recorded.

A flask, containing about 1000 cc. of sea-water treated with Miquel's solutions, was inoculated with approximately equal amounts of certain persistent cultures of diatoms, which we possessed at the time. The following diatoms were in this way introduced:—*Chaetoceras constrictum*, *Biddulphia mobiliensis*, *Skeletonema costatum*, *Coscinodiscus excentricus*, *Streptotheca thamensis*. The flagellate (*Chilomonas* sp.) was also introduced, since it was present in the culture of *Coscinodiscus*. The experiment was started on August 26th, 1907. On September 6th (11 days) *Biddulphia*, *Coscinodiscus* and *Chaetoceras* were increasing rapidly and were very healthy. *Skeletonema* was not so good, and no *Streptotheca* was found.

On October 2nd (37 days) *Biddulphia* was numerous and healthy, *Coscinodiscus* was healthy but not so numerous. *Skeletonema* was poor, and *Chaetoceras* was not seen. Flagellates (*Chilomonas*) had become very numerous.

On October 31st (66 days) all the diatoms were in very poor condition, *Coscinodiscus* being slightly better than the others. The flagellates (*Chilomonas*) were extremely thick, giving the water a deep red colour.

Subsequently a small green alga (*Pleurococcus mucosus*) appeared, having probably been derived from the *Coscinodiscus* culture. This increased very greatly in quantity, whilst the flagellates became inconspicuous.

On July 28th, 1909 (1 year 11 months), some *Coscinodiscus*, which were still in a healthy condition, were seen in a sample examined from the flask. A great quantity of *Pleurococcus*, in a healthy condition, was also present, but no other organisms were noted. On this date a subculture was made from the flask in normal outside Miquel. The subculture gave a considerable growth of *Skeletonema*, the cells being, however, of a very abnormal character, and a good many normal and

healthy *Coscinodiscus* were found in each sample examined. The whole culture was crowded with *Chilomonas* in a very active state, which gave the whole contents of the flask a deep red-brown colour. Up to August 24th, the green alga (*Pleurococcus*) had not become sufficiently abundant to be detected by the naked-eye appearance of the flask, though it could be seen in samples examined with the microscope.

In another experiment, a flask of Miquel sea-water was inoculated (May 4th, 1908) from two cultures, one containing the green alga (*Pleurococcus mucosus*) and the other *Thalassiosira decipiens*. At first both did well, and on May 20th (16 days) there was a very good crop both of the diatom and the alga. Gradually, however, the alga became predominant, and on October 14th (163 days) only quite empty frustules of *Thalassiosira* could be found, whilst the growth of *Pleurococcus* was abundant and healthy. The only case where a diatom was observed to flourish in the presence of this green alga was in a culture of *Nitzschia*, a bottom form. In this case a very abundant growth of the diatom was obtained, but the *Pleurococcus* did not multiply to any extent, although it could always be found on microscopic examination.

III. NOTES ON PARTICULAR SPECIES OF DIATOMS, ON THEIR METHODS OF REPRODUCTION, AND ON OTHER ALGÆ OCCURRING IN CULTURES.

A list has been already given (p. 425) of those species of diatoms which we have obtained in "persistent" cultures. Of these, a species belonging to the genus *Thalassiosira* has been used for experimental work in the great majority of cases. We are not quite certain as to the identity of the species, but since it most resembles *T. decipiens*, we have called it by that name, although it does not exactly conform to the published descriptions of that form. The most characteristic feature of this particular species is the eccentric marking on the valves, which are also seen on the valves of the diatom *coscinodiscus eccentricus*, Ehr., and, as is typical of the genus, the valves are united into chains by a delicate filament. Jörgensen (50, 51) describes the valves as "decidedly convex," Gran (49) as "flat," both agree that there are marginal spines and a single asymmetrical spine. Our cultural forms are united together by a filament into chains, some of which are made up of five hundred cells and more, but the distance between each is considerably smaller than that figured by Jörgensen. The valves are quite flat and the marginal spines are often

present, although this is not always the case. The apiculus can nearly always be seen, though it has not been observed in a few isolated instances. In one culture these markings were also easily seen or developed in it, but in none of the others started from this one have we seen any at all. The genus seems to be able to tolerate the conflicting descriptions due to variations in what is really the same species.

Persistent cultures of *Coscinodiscus* have been obtained, and it is interesting to note that they are rather exceptional as those commonly found with *Thalassiosira*. The filament joining the valves is broken. The two species are easily discriminated by a practical test.

Two species of the genus *Biddulphia* are present in our cultures, namely *Biddulphia regia*, M. Schultze, and *B. patula*, M. Schultze, as one species, but Ostensund (52) considers them really distinct. We have obtained them from several different sources, and they are easily recognizable, never confused with other diatoms. The colonies can be easily seen with the lens.

The most generally accepted account of the rigid character of the diatom is briefly that the valves undergo a decrease in size of the inner part during the life inside the old valve. So that the new valves are smaller as multiplication proceeds, which the original size can be seen in the formation of what are known as megafrustules, and these are the minimum limit of size having been reached. Special processes of reproduction have been noted in our work (p. 425).

The diatoms in our cultures are very different from those there is, in nearly every respect, compared with specimens

as were found in each sample examined. The whole of with *Thalassiosira* is a very active state, which gives it of the shell a deep red-brown colour. It is not seen (Microscopic) but not become sufficiently by the naked-eye appearance of the shell, as in samples examined with the microscope. Next, a flask of Miquel sea water was inoculated on two cultures, one containing the green alga and the other *Thalassiosira* *desipiens*. At first May 26th (16 days) there was a very good crop of the alga. Gradually, however, the alga became smaller. With (105 days) only quite empty ones could be found, while the growth of *Thalassiosira* was healthy. The only case where a diatom is in the presence of this green alga was in a flask from. In this case a very abundant one obtained, but the *Thalassiosira* did not though it could always be found on microscopic

ARTIFICIAL CULTURES OF MARINE PLANKTONIC ORGANISMS OF REPRODUCTION, AND OCCURRING IN CULTURES.

from (p. 425) of these species of diatoms "persistent" cultures. Of these, a species *Thalassiosira* has been used for experimental cases. We are not quite certain as to whether it most resembles *T. desipiens*, but none, although it does not exactly resemble that form. The most peculiar species is the concentric markings seen on the valves of the diatom and, as is typical of the genus, the by a delicate filament. Simpson (50), "eccentrically convex," Goun (49) as "lat," "girded spines and a single symmetrical united together by a filament into of five hundred cells and more, but hardly smaller than that figured by and the marginal spines are often

present, although this is not always the case. The odd segments spinous can nearly always be seen. The concentric markings have been observed in a few isolated cases, and are then usually very distinct. In one culture these markings on the valves were very distinct and were also easily seen on the megastrophules (cf. below) which developed in it, but in none of the several generations of cultures started from this one have we been able to find any traces of them at all. The genus seems to be in considerable confusion, and it is probable that the conflicting descriptions given by different observers are variations in what is really one species.

Persistent cultures of *Thalassiosira* *convexiuscula*, Ehrh., have also obtained, and it is interesting to note that this diatom sometimes chains, but they are rather exceptional. These chains are never as those commonly found with *Thalassiosira*, two or four cells and the rule. The filament joining the valves is also finer and more broken. The two species are quite distinct, and cultures of it be discriminated by a practised eye.

Two species of the genus *Balliophilus* are commonly met in our cultures, namely *Balliophilus* *multicellulosa* (Ball), Goun., and *Balliophilus* *oxya*, M. Schultze. These two forms are generally as one species, but Owsenfield (54) has recently shown that they are really distinct. We have obtained persistent cultures of both from several different samples of plankton, and the two are easily recognizable, never merging into one another. W. diatoms, inoculated from plankton (see p. 425), contain but the colonies can be easily distinguished with a microscope.

The most generally accepted theory of the reproduction of diatoms is briefly that the cells divide by simple fission, and, because of the rigid character of the cell walls each division results in a decrease in size of the new valves, since this must always occur inside the old valve. So the frustules gradually get smaller as multiplication proceeds, thus necessitating some process by which the original size can be re-established. This takes place by the formation of what are known as auxospores, which attain a minimum limit of size and then multiply by division into megastrophules, and these in turn multiply by division until a minimum limit of size has again been reached. There are several processes of reproduction, but no occurrence of any has been noted in our work (cf. Miquel, 54).

The diatoms in our cultures multiply by simple fission, and there is, in nearly every case, a considerable diminution in size compared with specimens from the plankton, this dim-

seems to reach a limit, where further decrease does not take place. In chains of *Thalassiosira*, several hundred cells in length, no difference in size between individuals could be made out. Auxospores are commonly formed with every species, but only in the cultures of *Coscinodiscus* and *Thalassiosira* have megafrustules been found, and in these they are very exceptional. These megafrustules seem to divide once or twice and then die or form new auxospores. What exactly is the fate of these auxospores, which are often exceedingly numerous, we have not been able to make out. It seems that cultural conditions are not favourable to this mode of reproduction, and that the auxospores do not further the multiplication of the diatom at all. If this were not the case, stages of the formation of auxospores into frustules must have been seen in some at least of the very numerous samples examined. As it is, what has been said to take place is, that the cell contents expand and force apart the valves of the diatom and emerge as a spherical body about three or four times the diameter of the parent cell. The chromatophores and diatomin then collect to one side, forming a compact cap against the cell wall. Beyond this point no stages have been found, except in the case of the few cultures where megafrustules were formed. In these the chromatophores, etc., gradually formed into the shape of the diatom (*Coscinodiscus*), the silicious coat with plain eccentric markings was easily seen inside the spore, and lastly, the cell wall of the spore burst, leaving the megafrustule free. The megafrustule was measured and found to have a diameter three times that of the parent cell.

In the case of the diatom we have very largely used for feeding larvae, etc., namely *Nitzschia closterium*, *forma minutissima*, a great number of cultures have been made, all originating from the single drop from which the first persistent culture was obtained. The total amount of growth in all the various cultures has been enormous, and the number of generations must be quite inconceivable. No diminution in size has, however, been appreciable, and no sign of any method of re-establishment of size has been seen, although these cultures have been under constant observation for over two years. This seems to prove that the theory of gradual decrease in size with successive generations cannot be generally applied.

The following experiment on the rate of multiplication of *Thalassiosira* in normal Miquel sea-water was carried out. A single drop from a fresh and vigorous culture was kept under a microscope as a hanging-drop preparation in a moist chamber. The number of diatoms in this drop was counted from time to time, and the results are given in the following table:—

Day.
11th
14th
19th
27th
34th
41st

The curve obtained number of days at progression, where the days. To show the intervals as the diatoms be seen that, after set in, the number that every diatom in normal cultural this figure on account

Besides diatom We are indebted unicellular alga, attempting to obtain *Pleurococcus mucronatus* introduced into the expense of hardy and cultured definitely. Much occur, but the effective reproduction

In cultures in and filamentous *Rhodophyceae* have been able to form *Chlorophyceae* are interesting to the plants of *Laminaria*. Mr. Drew (4) has its early life in would without rich material.

Many forms Among these obtained in

Day.	Number of frustules.	Geometric progression.
11th	59	63
14th	62	68
19th	85	85
27th	140	120
34th	170	160
41st	190	220

The curve obtained by plotting the number of diatoms against the number of days approximates the curve of an ordinary geometric progression, where the ratio is two and the periods are equal to sixteen days. To show this the figures read off from the curve at the same intervals as the diatoms are appended in the table. From this it will be seen that, after a start had been made and before exhaustion set in, the numbers obtained agree fairly closely with the assumption that every diatom divided once in a period of sixteen days. Probably in normal cultural conditions the rate of multiplication greatly exceeds this figure on account of better lighting, etc. (cp. Miquel, 12).

Besides diatoms, many other organisms appear in these cultures. We are indebted to Prof. G. S. West for the identification of a form of unicellular alga, which is very common and difficult to avoid when attempting to obtain persistent cultures of the *Diatomaceae*, namely *Pleurococcus mucosus* (Kutz.), Rabenh. This small green alga, if once introduced into a culture of a plankton diatom, will soon multiply at the expense of the latter with its ultimate extinction. It is very hardy and cultures of it in almost every medium seem to last indefinitely. Multiplication beyond a certain point probably does not occur, but the cells retain their colour and normal shape and will start active reproduction if suitable nutrient material is provided.

In cultures inoculated from plankton many other forms of unicellular and filamentous algae thrive. Several species belonging to the classes *Rhodophyceae* and *Myxophyceae* commonly occur, but we have not been able to identify them. The most usual filamentous forms of *Chlorophyceae* are *Enteromorpha*, *Vaucheria*, *Rhizoclonium*, etc. It is interesting to note that it was the unintentional appearance of young plants of *Laminaria digitata* in some of our Petri dishes that led Mr. Drew (4) to cultivate this alga in Miquel sea-water and so discover its early life history. Cultivations of marine algae by these methods would without doubt yield many new species, and would also provide rich material for the study of their modes of reproduction.

Many forms of flagellates live either together with diatoms or alone. Among these is an unidentified species of *Chilomonas* which we have obtained in persistent culture. It multiplies very rapidly, colouring

the whole medium a deep red-brown. It flourishes in Miquel sea-water, and its nutrition is evidently autotrophic. In one culture in Miquel sea-water inoculated with plankton a number of *Coccospheres* developed, probably *Coccosphaera atlantica*, Ostenf. Other flagellates and ciliated infusoria are very commonly met with, such as *Bodo*, *Euplotes*, *Euglena*, etc., which all seem to depend on the diatoms or other vegetable organisms for their food material.

IV. THE REARING OF MARINE LARVÆ.

In the rearing of pelagic larval forms of marine animals,* the principle which we have followed has been to introduce into pure, sterile sea-water the larvæ to be reared, together with a pure culture of a suitable food. As far as practicable all other organisms have been excluded from the rearing vessels. It should be added that the food used in all successful experiments has been of a vegetable nature, and has continued to grow actively in the vessels. This is important from the point of view of oxygen supply. Under the above conditions, or rather under the nearest approach to them at which we have been able to arrive, no change of water has been found necessary.

Methods.—It will, perhaps, best make the matter plain if we first of all describe the actual procedure which we now follow in the case of such an animal as *Echinus esculentus* or *E. acutus*. The water to be used is first of all prepared by treating water from the aquarium tanks with powdered animal charcoal, filtering it through a Berkefeld filter (p. 431), and collecting it in sterilized glass vessels. All instruments and pipettes are sterilized by baking in an oven, and a fresh sterile pipette is used for each operation during the progress of the work. Specimens of *Echinus* are then opened until a perfectly ripe female has been found, that is to say, one in which the eggs separate quite freely, when a portion of the ovary is shaken in sea-water.

Pieces of ovary taken from a little below the exposed surface are then placed in sterile sea-water in a shallow glass dish and shaken with forceps in order to get the eggs well separated, or a number of eggs from the centre of the ovary are drawn up with a pipette and placed in the water. A very small quantity of active sperm from a ripe male is then added, very little being sufficient to fertilize a large number of eggs. Excess of sperm should be avoided owing to its liability to putrefy. After an interval of ten or fifteen minutes the water containing the eggs is filtered through bolting silk of 100 meshes per inch, which just allows single eggs to pass through, whilst keeping

* See Bibliography; especially Grave (26), MacBride (28-30), Doncaster (25), etc.

back clusters of eggs amongst a number of water, and the beakers placed where the temperature is above 15° C. In the vessels will swim up to the surface of this shape. to jars* of sterile sea-water each jar of 2000 c.c. : a pure culture of *Nitzschia closterium*, its size is suitable, floating throughout mixture with the larvæ even a temperature the larvæ have met with six to nine weeks time to time and diatoms do not grow added from the culture towards the end of diatoms. In this some of the water is sterile sea-water. food at the beginning able to feed as soon

The method just described is of no detriment to the eggs effected by heat. The water may be aerated by vigorous shaking of "tank-water," in the ordinary way, to

With regard to a variety of the eggs of a number of other organisms experimented. sometimes be reared with a fine-mesh

* The vessels were of 2000 c.c., which was filled with water in which the cork balls

† In hot weather the water, which may

TON ORGANISMS.

Miquel sea-water,
culture in Miquel
Coccospheres de-
Other flagellates
h, such as *Bodo*,
the diatoms or

VÆ.

animals,* the
duce into pure,
a pure culture
isms have been
that the food
ole nature, and
important from
conditions, or
have been able

if we first of
in the case
e water to be
uarium tanks
rkefeld filter
instruments
fresh sterile
of the work.
ripe female
parate quite

surface are
and shaken
a number
pipette and
perm from
fertilize a
l owing to
minutes the
00 meshes
et keeping

ON THE ARTIFICIAL CULTURE OF MARINE PLANKTON ORGANISMS. 465

back clusters of eggs or other large material. The filtrate is divided amongst a number of tall narrow beakers containing sterile sea-water, and the beakers, after being covered with a glass plate, are placed where the temperature will be uniform and not rise much above 15° C. In the course of twenty-four hours the healthy larvæ will swim up to the surface and can be easily seen and removed from vessels of this shape. They are transferred by means of sterile pipettes to jars* of sterile sea-water, about fifty to seventy larvæ being put in each jar of 2000 c.c. sea-water. At the same time, a good pipetteful of a pure culture of diatom is added to each jar. The small diatom *Nitzschia closterium*, *forma minutissima*, we have found most useful, as its size is suitable, and it grows well in animal-charcoal tank-water, floating throughout the body of the water, and so being in intimate admixture with the larvæ. The jars are placed in a moderate light and at as even a temperature as possible.† No further attention is necessary until the larvæ have metamorphosed. The metamorphosis takes place in from six to nine weeks after fertilization. Larvæ may be taken out from time to time and examined to see if they are feeding well. If the diatoms do not grow sufficiently rapidly in the jar, more should be added from the culture flasks. We are more often troubled, however, towards the end of an experiment by an excessive abundance of diatoms. In this case the jar may either be put in a darker place or some of the water may be drawn off and replaced by a fresh supply of sterile sea-water. Care should be taken to have a sufficient supply of food at the beginning of the experiment, so that the larvæ may be able to feed as soon as they are ready for food.

The method just described can be modified in various ways without detriment to the result. Sufficient sterilization of the water may be effected by heating to 70° C. for fifteen minutes, after which it should be aerated by violent shaking; "outside" water may be used instead of "tank-water," and may be treated with Miquel's solutions in the ordinary way, to ensure a satisfactory growth of the food-diatom.

With regard to the food organisms, we have tried to obtain as large a variety of these in pure culture as possible, and then to make trial of a number of them with each batch of larvæ on which we have experimented. If no suitable pure cultures are available, success can sometimes be obtained by adding a few drops of tow-netting, collected with a fine-meshed net (180 meshes per inch), directly to the treat-

* The vessels we use are ordinary green glass sweet-jars, having a capacity of about 2000 c.c., which are kept covered with the glass stoppers provided with such jars, from which the cork band has been removed.

sterile water containing the larvæ. In this case one depends on the chance of a suitable food organism growing in the vessel unaccompanied by any destructive organism. On several occasions a satisfactory result has been reached by proceeding in this way, and the method is generally worth a trial, seeing that the number of larvæ obtainable from an ordinary fertilization is very large, and many different experiments are easily made with them.

We will now give details of some of the results obtained by making use of the methods described, or of their modifications.

Echinus acutus.—The first successful experiment was made with this species. Eggs fertilized on June 13th, 1905, produced healthy larvæ, 50 to 75 of which were placed, three days later, in a glass jar containing 2000 c.c. of outside sea-water, filtered through animal charcoal, to which modified Miquel solutions were added. They were fed on a diatom culture, containing a small species of *Chaetoceras*, which did not form chains, a small diatom probably belonging to the genus *Melosira*, a small Naviculoid diatom, two minute flagellates, and a small green organism, probably one of the *Pleurococcaceæ*. The vessel stood in a shallow tank, through which a stream of aquarium water was flowing, and the temperature was fairly constant at 15° or 16° C., though there is one record of 19° C. at the end of July. The first two young *Echinus* were seen on July 25th, 42 days after fertilization, and on August 1st 20 were counted. On August 5th (the 53rd day) a careful search through the jar gave 21 young *Echinus* of normal size attached to the glass, 6 minute but fully formed *Echinus*, about 23 still in the *Pluteus* stage, roughly half of which were well advanced. On August 16th some of the water, which had not been changed since the beginning of the experiment, was replaced by "outside" water. On October 5th (16 weeks after fertilization) 12 *Echinus* were still alive. Some pieces of red seaweed were placed in the jar, upon which the *Echinus* fixed themselves and fed. Several of these specimens lived for over a year, but sufficient attention was not given to finding suitable food for them after the metamorphosis, so that they did not grow very large.

Echinus esculentus.—Three successful experiments have been made with *E. esculentus*. In the first (eggs fertilized April 5th, 1907) "outside" water treated with animal charcoal and filtered through filter-cloth, but not otherwise sterilized, was used. A number of jars of 2000 c.c. capacity containing larvæ were set up, and, to the most of these, various diatom cultures then in our possession were added, none of which, however, gave a satisfactory result. In two jars, on the

other hand, to which growth of diatoms and first young *Echinus* but may have been to 40 metamorpho temperature varied

In the second ex similar water, the *closterium*, var., an (48 days after fer The temperature

In the third ex tank-water treat a Berkefeld filter flagellate (probab *Echinus* were fou probably metam *Nitzschia closteri* because the gro experiment.

Echinus mili charcoal Berke In one jar the fed on a pure thirty-eight d phosed. On climbing abov January 8th, 16" dropping 15° to 10° C.

To anothe fresh Plank flagellates a phosed.

In a seco the larvæ v few well-a pleted the

Cucuma containing segmented

other hand, to which no culture was added, there was considerable growth of diatoms and of a flagellate, upon which the Plutei fed. The first young *Echinus* were recorded in both jars on June 8th (64 days), but may have been present a few days earlier. Eventually from 30 to 40 metamorphosed in one jar and about 12 in the other. The temperature varied from 10.5° C. to 12.5° C.

In the second experiment (eggs fertilized June 8th, 1908), made with similar water, the larvæ were fed on a pure culture of *Nitzschia closterium*, var., and six had completely metamorphosed on July 26th (48 days after fertilization), two more subsequently coming through. The temperature was generally 15° to 16° or 17° C.

In the third experiment (eggs fertilized March 29th, 1909) aquarium tank-water treated with animal charcoal and then filtered through a Berkefeld filter was used. Plutei, fed with a pure culture of a small flagellate (probably *Chilomonas* sp.) grew satisfactorily, and eight young *Echinus* were found on June 5th (68 days after fertilization), which had probably metamorphosed some days earlier. Two other jars in which *Nitzschia closterium*, var., was used as food, were not successful, probably because the growth of diatoms became too thick towards the end of the experiment.

Echinus miliaris. In the first experiment with this species animal-charcoal Berkefeld water was used, each jar containing as usual 2000 c.c. In one jar the Plutei, from eggs fertilized on August 27th, 1907, were fed on a pure culture of *Nitzschia closterium*, var. On October 4th, i.e. thirty-eight days after fertilization, one *Echinus* had just metamorphosed. On October 29th about a dozen healthy-looking *Echini* were climbing about the jar, and many were still in a healthy condition on January 8th, 1908. Temperatures: September, 15° to 19° C., October, 16° dropping to 13° towards end, November, 12° to 11° C., December, 15° to 10° C.

To another jar containing larvæ from the same batch a few drops of fresh Plankton were added as food. The Plutei in this case fed on flagellates and *Nitzschia* which grew in the jar, and several metamorphosed.

In a second experiment with eggs fertilized on September 13th, 1907, the larvæ were fed with *Nitzschia closterium*, but although there were a few well-advanced Plutei still living on January 8th, 1908, none completed the metamorphosis.

Cucumaria saxicola. A female *Cucumaria*, one of a number in a dish containing "outside" water, laid eggs, which were fertilized, and segmented on May 12th, 1906. A number of these were placed in a

flask in 800 c.c. of "outside" water, which had been sterilized by heating and then treated with animal charcoal and filtered. About 1 c.c. of fine plankton, containing diatoms, was added to the flask on May 12th. On May 25th some of the water was poured off and a new supply added. As the amount of food seemed small, some culture of a green alga (*Pleurococcus mucosus* (Kütz.), Rabenh.) was added, and this continued to grow well in the flask. The larvæ continued healthy and formed young *Cucumaria*, of which many were still alive on July 25th, 1907, i.e. fourteen months after fertilization. Some of the water was changed in this flask on May 30th, 1906, June 18th, 1906, September 15th, 1906, and July 25th, 1907. Although many of these *Cucumaria* remained quite healthy they did not grow to any great size. Probably the food which was suitable to the larvæ and early stages, ought to have been changed as the animals grew older.

Pomatoceros triqueter. The larvæ of *Pomatoceros* are perhaps the easiest to rear, and give the most certain results of any with which we have experimented. They do well on the minute variety of *Nitzschia closterium*, but will feed upon almost any small diatom. Since the adults live in calcareous tubes attached to stones, and the tubes have to be broken open before the eggs can be obtained, it is not easy to get the latter free from infection of other organisms. If, therefore, the eggs are fertilized and placed in sterilized animal-charcoal water with only moderate precautions, sufficient growth of diatoms or other organisms will generally take place in the jar to feed the larvæ and bring them to the adult state. When once fixed to the glass the worms are very hardy and healthy, and a stream of ordinary aquarium water can be run through the jar. They then grow rapidly and attain a size equal to any found on the shore. The following experiment may be given in detail to illustrate the time occupied in development. On August 29th, 1907, eggs of *Pomatoceros triqueter* were fertilized in animal-charcoal Berkefeld water, and some pure culture of *Nitzschia closterium*, var., added. The larvæ fed well, and on October 1st (i.e. thirty-three days after fertilization) a great number had fixed on the sides of the jar and made quite normal tubes. A constant stream of the ordinary aquarium water was then allowed to run through the jar, and the worms continued to grow and flourish, reaching a large size, and are still alive and healthy (November, 1909). A similar result was obtained from the same batch of eggs by feeding on a pure culture of a flagellate infusorian. Temperatures during these two experiments were between 15° C. and 19° C.

Chaetopterus variopedatus. Four experiments were made with this

species. The food was *Nitzschia closterium* 1908, fed on this in an advanced stage. They formed tubes. Two using flagellates as food.

Sabellaria alveolaris species on eggs fed in "outside" water containing animal charcoal on a pure culture of *Nitzschia closterium* and developed well. Eaten with a sucrose solution to 12° and 9° C., in three days were hatched. The temperature from 15° to 19° C. and continued a month in October.

Archidoris tubicola. The larvæ of *Archidoris* have been successfully reared. Larvæ of *Archidoris* hatched out on August 15th 1908. Collected from the plankton was added to another flask. Culture of the *Nitzschia closterium* the larvæ in the green alga. They were active hatching, and on August 15th 1908 on the bottom of decomposition tissue looked like seen. The green *Pleurococcus* appeared mucous.

species. The food which gave most promise of success was the diatom *Nitzschia closterium*, var. Larvæ from eggs fertilized on July 20th, 1908, fed on this material lived until October 30th, and reached an advanced stage. They did not, however, adopt the adult habit and form tubes. Two larvæ were also reared to an advanced stage by using flagellates and, in later stages, the diatom *Skeletonema costatum* as food.

Sabellaria alveolata. One experiment only was made with this species on eggs fertilized on July 19th, 1908. The eggs were fertilized in "outside" water and the larvæ subsequently transferred to jars containing animal-charcoal Berkefeld aquarium water. They were fed on a pure culture of *Nitzschia closterium*, var., and kept healthy and active and developed well until nearly the end of October, when, simultaneously with a sudden drop in the temperature from 15° and 16° C. to 12° and 9° C., they sank to the bottom of the vessel and in about three days were all dead. Temperatures:—During July and August, the temperature kept fairly constant at about 17° C., with a range from 15° to 19° C. During September it was generally about 15° C., and continued at about this level until the fall in the middle of October.

Archidoris tuberculata. A good many trials have been made to rear the larvæ of nudibranchiate molluscs, but up to the present not much success has been achieved. The best experiment was one made with larvæ of *Archidoris tuberculata*. A number of veligers of this species hatched out on May 8th, 1908, from some spawn, which had just been collected from the shore. Some of these were put in a flask containing 1000 c.c. of sterilized animal-charcoal water and about 1 c.c. of fine plankton was added. On May 14th a few veligers were transferred to another flask of sterilized animal-charcoal water and some pure culture of the green alga, *Pleurococcus mucosus*, was added. Whereas the larvæ in the original flask did not live long, those provided with the green alga fed well and developed for some time. A number of them were active and vigorous on July 4th, i.e. 51 days after hatching, and several were still swimming at the end of July. On August 15th none could be seen moving, but two of those which lay on the bottom, when examined with the microscope, showed no sign of decomposition. The animal was retracted in the shell, but the tissue looked healthy, and the eye-spots and otoliths could be seen. The growth in the flask seemed to be a quite pure culture of *Pleurococcus*. Larvæ were examined again on September 14th, and appeared much as in August, the tissue still showing no sign of dis-

integration. The flask was not again examined microscopically until July 25th of the following year (1907). No sign of the larvæ could then be seen, but the culture of *Pleurococcus* remained pure and healthy.

Subsequent experiments were made with spawn, which was deposited by the females in confinement. Although the spawn hatched and gave apparently healthy larvæ, these did not live for more than a few days.

Calanus finmarchicus. A single experiment is perhaps worth recording, as showing that it ought to be possible to rear this species without great difficulty. On August 8th, 1905, to a flask containing 1000 c.c. of outside water (unsterilized) there was added $\frac{1}{2}$ c.c. of Miquel's solution B and $\frac{1}{2}$ c.c. of a 1.5 per cent solution of anhydrous sodium carbonate. A few *Calanus finmarchicus* and some decapod Zoeas were put in, together with a quantity of a culture containing mixed diatoms. On September 8th all the Zoeas were dead, but three *Calanus* were alive, and *Nitzschia* and a number of bottom diatoms were very plentiful. On September 17th the three large *Calanus* were alive and vigorous, and a considerable number of *Nauplii* were seen in the flask. By September 22nd two of the *Nauplii* had developed into young *Calanus*. These, however, did not live for more than a week or ten days, and the adults also died. The flask was abandoned on November 13th, the water in it not having been changed since the commencement of the experiment.

Echinus hybrid. A successful experiment on crossing *E. esculentus* and *E. acutus* was carried out by Mr. W. De Morgan, who was working at the Plymouth Laboratory. We provided him with treated water and diatom cultures for food, and he followed our methods. We are indebted to him for allowing us to publish these results. Some eggs from a ripe *E. esculentus* were fertilized by active sperm from *E. acutus* in sterilized water on March 29th, 1900. Healthy larvæ were obtained and were transferred two days later to tank-water, which had been treated with animal charcoal and filtered through a Berkefeld filter. A culture of *Nitzschia closterium*, var., was added as food, and the larvæ developed rapidly, feeding well. Several were completely metamorphosed on May 7th, or thirty-nine days after fertilization. In all thirty young hybrids were obtained, and a number of these are still alive and feeding on red weeds.

Sacculina carcini. Mr. Geoffrey Smith has recorded the fact (*Quart. Journ. Micr. Sci.*, LI, 1907, p. 625) that he was able to rear the larvæ of *Sacculina* up to the Cypris stage, when they attached themselves to

their host, *Carcinus maenas*. The tank-water treated with animal charcoal and passed through a Berkefeld filter. In this case the larvæ do not feed after hatching, but that these larvæ had previously

Summary of Method for Rearing Marine Plankton.—The best results in rearing marine plankton require the following precautions:—

1. The eggs of the female and spermatozoa of the male should be used.
2. The smallest quantity of food organisms should be used.
3. Sterile sea-water, treated with animal charcoal and filtered through a Berkefeld filter, should grow well in it, should be used.
4. All dishes, jars, instruments, etc., should be sterilized before use, to prevent the introduction of other organisms than the larvæ to be reared. The jars should be covered.
5. The eggs after fertilization should be kept in a solution of foreign matter, pieces of seaweed, etc., so that the larvæ swim up they may feed on it.
6. The food organisms should be distributed in such a way that they can draw them into their mouths. They should distribute its food so that it does not settle too readily on the bottom. This is the great advantage of using *minutissima*.)
7. The food should be changed frequently, but should commence feeding on it as soon as possible, however, must not be changed too often.
8. The temperature should be kept within the limits, the actual temperature should be the avoidance of it.
9. A good north light should be used for the rearing.

their host, *Carcinus maenas*. These larvæ were kept in aquarium tank-water treated with animal charcoal and filtered through a Berkefeld filter. In this case the question of food did not arise, as the larvæ do not feed after hatching. It must be noted, however, that these larvæ had previously been reared by Müller and by Delage.

Summary of Method for Rearing Larvæ. We have found that the best results in rearing marine larvæ have been attained by taking the following precautions:—

1. The eggs of the female selected must be really ripe, and the spermatozoa of the male active.
2. The smallest quantity of sperm necessary to fertilize the eggs should be used.
3. Sterile sea-water, treated in such a way that diatoms, etc., will grow well in it, should be used. No frequent change of water is then necessary.
4. All dishes, jars, instruments, and pipettes should be carefully sterilized before use. Every possible effort should be made to prevent the introduction into the rearing-jars of any organisms other than the larvæ to be reared, and organisms on which they feed. The jars should be covered with loosely fitting glass covers.
5. The eggs after fertilization must be separated from all foreign matter, pieces of ovary, or testis, etc. As soon as the larvæ swim up they should be pipetted off into fresh vessels of treated water, so as to leave behind any unsegmented eggs, etc.
6. The food organisms should be small in size, so that the larvæ can draw them into the mouth by ciliary currents. The food should distribute itself through the body of the liquid and not settle too readily on the bottom of the vessel. (This is one of the great advantages of the diatom *Nitzschia closterium*, *forma minutissima*.)
7. The food should be abundant early, so that the larvæ may commence feeding as soon as they are able to do so. The food, however, must not be allowed to get excessively thick in the water. It can be kept down by diminishing the light or by changing some of the water.
8. The temperature should be kept as constant as possible. Within limits, the actual degree of temperature is not so important as the avoidance of rapid change of temperature.
9. A good north light, not exposed to direct sunlight, is most suitable for the rearing-jars.

10. In determining the amount of water to be used in any particular vessel, regard must be had to the amount of water surface exposed to the air, which should be large in proportion to the volume of the water.
11. A change of food is generally required after the metamorphosis of the larvæ.

BIBLIOGRAPHY.

Cultures.

1. Beijerinck, M. W.—Das Assimilationsprodukt der Kolensäure in den Chromatophoren der Diatomeen. Rec. Trav. Bot. Neerland, I, 1904, p. 28.
2. Benecke, W.—Über farblose Diatomeen der Kieler Förde. Pringsh. Jahrb., XXXV, 1900, p. 567.
3. Breazeale, J. F.—Effect of Certain Solids upon the Growth of Seedlings in Water Cultures. Botanical Gazette, XLI, 1906, p. 54.
4. Drew, G. H.—The Reproduction and early Development of *Laminaria digitata* and *L. saccharina*. Annals of Botany, XXIV, 1910, p. 177.
5. Gill, C. Houghton.—See Van Heurck, H. 21 and 57.
6. Karsten, G.—Die Formänderungen von *Skeletonema costatum*, Grun., und ihre Abhängigkeit von äusseren Faktoren. Wiss. Meeresunters. Kiel. N.F., III, 1898, p. 5.
7. Karsten, G.—Ueber farblose Diatomeen. Flora, LXXXIX, 1901, p. 404.
8. Lockwood, S. M.—Raising Diatoms in the Laboratory. Journ. New York Micr. Soc., 1886, p. 153.
9. Lockwood, S. M.—Aberrant Forms in cultivated Diatoms. Amer. Monthly Micr. Journ., 1893, p. 259.
10. Lockwood, S. M.—Forms anormales chez les Diatomées cultivées artificiellement. Le Diatomiste, II, 1893-96, p. 9. Ann. de Micrographie, X, 1898, p. 1.
11. Miquel, P.—De la Culture artificielle des Diatomées. Le Diatomiste, I, 1890-93, pp. 73, 93, 121, 149, 165. Le Micrographie Préparature, V, 1897, p. 69.
12. Miquel, P.—Recherches expérimentales sur la physiologie, la morphologie, et la pathologie des Diatomées. Ann. de Micrographie, IV, 1891-92, pp. 273, 321, 408, 520; V, 1893, pp. 437, 521; X, 1898, pp. 49, 177, 182. Le Micro. Préparature, XI, 1903, p. 174; XII, 1904, p. 32.
13. Miquel, P.—Du rétablissement de la taille et de la rectification de la forme chez les Diatomées. Le Diatomiste, II, 1893-96, pp. 61, 88.
14. Miquel, P.—Des Spores des Diatomées. Le Diatomiste, II, 1893-96, p. 26.
15. Miquel, P.—Du noyau chez les Diatomées. Le Diatomiste, II, 1893-96, p. 105. Le Micro. Préparature, XII, 1904, p. 167; XIII, 1905, p. 83.
16. Richter, O.—Reinkulturen von Diatomeen. Ber. deut. bot. Gesell., XXI, 1903, p. 493.
17. Richter, O.—Über Reinkulturen von Diatomeen und die Notwendigkeit der Kieselsäure für *Nitzschia palea* (Kütz.). W. Sm. Verh. d. Gesell. deut. Naturf. u. Äzzte. Breslau, II, 1904, p. 249.
18. Richter, O.—Zur physiologie der Diatomeen. S. B. K. Akad. Wiss. Wien, CXV, 1906, p. 935.
19. Richter, O.—Ueber die Notwendigkeit des Natriums für eine farblose Meeresdiatomee. Wiener-Festschrift. Wien, 1908, p. 167.
20. Senft, E.—Ueber die Agar-Agar Diatomeen. Zeit. d. Allgem. öst. Apotheker-Vereines, 1902, n. 9, figs. 1-9.

21. Van Heurck, H.—Notice b 1893-96, p. 125.
22. Van Heurck, H.—Culture pp. 195 and 225.
23. Agassiz, A.—Revision of
24. Cowles, R. F. Notes on Hopkins Univ.
25. Doncaster, L.—On rearing Phil. Soc., XII, p. 579.
26. Grave, C.—A Method
27. Lillie, R. S.—The Stru Mittheil. Zool.
28. MacBride, E. W.—The Dover, 1899
29. MacBride, E. W.—No Biol. Assoc., I
30. MacBride, E. W.—The points in the Roy. Soc., B.
31. Thecl, H.—On the De Sci., Upsala,
32. Zeleny, C.—The Rea the Young A
33. Dittmar, W.—On the 1873-76, CH
34. Fox, C. J. J.—On the Distilled w Circumstanc Copenhagen
35. van 't Hoff, J. H.—2 weig, 1905.
36. Knudsen, M.—Hydr Copenhagen
37. Krogh, A.—On the in the sea. hagen.
38. Raben, E.—Über qu wasser.
39. Raben, E.—Quanti Wissensch
40. Ringer, W. E.—D v.h. onde
41. Ringer, W. E., and verbindv zoek d. 2
42. Salm, E.—Studie p. 471.
43. Tornøe, H.—On Expdt.

ON THE ARTIFICIAL CULTURE OF MARINE PLANKTON ORGANISMS. 473

21. Van Heurck, H.—Notice biographique sur C. Houghton Gill. *Le Diatomiste*, II, 1893-96, p. 125.
22. Van Heurck, H.—Culture des Diatomées. *Zeit. f. angew. Mikrosk.*, III, 1897, pp. 195 and 225.

Rearing.

23. Agassiz, A.—Revision of the Echini. Cambridge, Mass., 1872-4.
24. Cowles, R. P. Notes on the Rearing of the larvæ of *Polygordius*. Johns Hopkins Univ. Circulars XXII, No. 161, 1903.
25. Doncaster, L.—On rearing the later stages of Echinoid Larvæ. Cambridge Phil. Soc., XII, 1903, p. 48.
26. Grave, C.—A Method of Rearing Marine Larvæ. *Science*, N.S. xv, 1902, p. 579.
27. Lillie, R. S.—The Structure and Development of the Nephridia of *Arenicola*. *Mittheil. Zool. Sta. Neapel*, XVII, 1904-6, p. 341.
28. MacBride, E. W.—The Rearing of Larvæ of Echinidæ. *Reports, Brit. Assoc.*, Dover, 1899, p. 438.
29. MacBride, E. W.—Notes on the Rearing of Echinoid Larvæ. *Journ. Mar. Biol. Assoc.*, N.S. VI, 1900-3, p. 94.
30. MacBride, E. W.—The Development of *Echinus esculentus*, together with some points in the Development of *E. miliaris* and *E. acutus*. *Phil. Trans. Roy. Soc.*, B. CXCv, 1903, p. 285.
31. Theél, H.—On the Development of *Echinocyamus pusillus*. *Nova Acta R. Soc. Sci.*, Upsala, 1892.
32. Zeleny, C.—The Rearing of Serpulid Larvæ, with Notes on the Behaviour of the Young Animals. *Biol. Bull. Woods Holl.*, VIII, 1905, p. 308.

Chemistry.

33. Dittmar, W.—On the Alkalinity of Ocean Water. *Rept. Challenger Expdt.*, 1873-76, Chem. I, p. 124. London, 1884.
34. Fox, C. J. J.—On the co-efficients of Absorption of the Atmospheric gases in Distilled water and Sea-water. Part II, "Carbonic acid." *Publ. de Circonstance. Conseil internat. pour l'expl. de la mer.* No. 44. 1909. Copenhagen.
35. van 't Hoff, J. H.—Zur Bildung der ozeanischen Salzablagerungen. *Braunschweig*, 1905.
36. Knudsen, M.—Hydrography. *Danish Ingolf-Expedtn.*, I, part 2, p. 21, 1899. Copenhagen.
37. Krogh, A.—On the tension of Carbonic Acid in natural waters, and especially in the sea. *Meddelelser om Grønland*, XXVI, 1904, p. 333. Copenhagen.
38. Raben, E.—Über quantitative Bestimmung von Stickstoffverbindungen in Meerwasser. *Wissensch. Meeresunters.* Kiel, VIII, 1905, pp. 81, 279.
39. Raben, E.—Quantitative Bestimmung der im Meerwasser gelösten Kieseläure. *Wissensch. Meeresunters.* Kiel, VIII, 1905, p. 99, 286.
40. Ringer, W. E.—Die Alkalinität des Meereswassers. *Verh. uit. Rijksinstituut v.h. onderzoek d. zee*, IV, 1909. Helder.
41. Ringer, W. E., and Klingen, F. I. M. P.—Über die Bestimmung von Stickstoffverbindungen im Meereswasser. *Verh. uit. Rijksinstituut v.h. onderzoek d. zee*, I, 1907-8. Helder.
42. Salm, E.—Studie über Indikatoren. *Zeitsch. f. Phys. Chemie.* LVII, 1906, p. 471.
43. Tornøe, H.—On the Carbonic acid in Sea-water. *Norwegian North-Atlantic Expdt.* (1876-8), Chem. II, p. 24. Christiania, 1880.

General.

44. Baldwin, H. B., and Whipple, G. C.—Observed relations between Oxygen, Carbonic acid, and Algae growths in Weequahie Lake, Newark, New Jersey. Papers and Repts. Amer. Public Health Assoc., xxxii, 1906.
45. Brandt, K.—Beiträge zur Kenntniss der chemischen Zusammensetzung des Planktons. Wissensch. Meeresunters. Kiel, iii, 1898, p. 43.
46. Brandt, K.—Ueber den Stoffwechsel im Meere. Wissensch. Meeresunters. Kiel, iv, 1899, p. 215; vi, 1902, p. 23.
47. Brandt, K.—Ueber die Bedeutung der Stickstoffverbindungen für die Produktion im Meere. Beihefte z. Bot. Centralblatt., xvi, 1904.
48. Brandt, K.—On the production and conditions of production in the sea. Rapports et Procès-Verbaux, Conseil internat. pour l'expl. de la mer, iii, Appdx. D. Copenhagen, 1905.
49. Gran, H. H.—Diatomeen. Nordisches Plankton. Kiel u. Leipzig, 1903. Botanischer Teil, xix.
50. Jørgensen, E.—The Protist plankton and diatoms in bottom samples. Hydrographical and Biological Investigations. Bergen, 1906.
51. Klebahn, H.—Ein Überblick über die neuere Diatomeenliteratur. Archiv. für Protistenkunde, i, 1902, p. 421.
52. Lemmermann-Bremen, E.—Flagellatae, Chlorophyceae, Coccosphaerales, und Silicoflagellatae. Nordisches Plankton, Kiel u. Leipzig, 1903. Botanischer Teil, xxi.
53. Oltmanns, F.—Morphologie und Biologie der Algen (with bibliography). Two vols. Jena, 1905.
54. Ostenfeld, C. H.—On the immigration of *Biddulphia sinensis*, Grev., and its occurrence in the North Sea during 1903-7, etc. Meddelelser fra Komm. for Havundersøgelser. Plankton Series 1, No. 6, Copenhagen, 1908.
55. Pfeffer, W.—The Physiology of Plants (English edition). Three vols. Oxford, 1900.
56. Smith, G. P. Darnell.—On the oxidation of Ammonia in Sea-water. Journ. Mar. Biol. Assoc., N.S. iii, 1893-5, p. 304.
57. Van Heurck, H.—A Treatise on the Diatomaceae (translated by W. E. Baxter). London, 1896.
58. Vernon, H. M.—The relations between Marine Animal and Vegetable Life. Mittheil. Zool. Sta. Neapel, xiii, 1893-9, p. 341.
59. Whipple, G. C.—Some experiments on the Growth of Diatoms. Technology Quarterly, ix, 1896, p. 145. Boston.
60. Whipple, G. C.—The Microscopy of Drinking Water (and bibliography). New York, 1908.
61. Whipple, G. C., and Jackson, D. D.—*Asterionella*, its Biography, its Chemistry, and its effect on water supplies. Journ. N.E. Waterworks Assn., xiv, 1899, p. 1.
62. Whipple, G. C., and Parker, H. N.—On the amount of oxygen and carbonic acid dissolved in natural waters, and the effect of these gases upon the occurrence of microscopic organisms. Trans. Amer. Micr. Soc., xiii, 1901-2, p. 103.

On the Species

As remarked by "Upogebia, Leach covered by the in (1808) as *Cancer* notice of writers genus was Upogel Leach's own alter

Some doubt st and *Gebia deltu* same species.

Leach gives ex table xxi, figs. 1- distinct species, *Gebia stellata*, w were discovered whether Mr. So

Bell (*British* and *Gebia deltu* and gives good they are distin it be indeed di ing particulars less than twic The carapace i legs are more as long as it thicker, and t external ante figure, not me broader, more almost meml voluminous t